Homework for October 29, 2017.

Algebra.

Review the classwork handout and complete the exercises. Solve the remaining problems from the previous homework. Solve the following problems (some of the problems are repeated from the previous homeworks, please skip the ones you've already solved).

- 1. Prove that the following number is irrational.
 - a. $\sqrt{3}$
 - b. $\sqrt{5}$
- 2. Using mathematical induction, prove that $\forall n \in \mathbb{N}$,

a.
$$\sum_{k=1}^{n} (2k-1)^2 = 1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{4n^3 - n}{3}$$

b.
$$\sum_{k=1}^{n} (2k)^2 = 2^2 + 4^2 + 6^2 + \dots + (2n)^2 = \frac{2n(2n+1)(n+1)}{3}$$

c.
$$\sum_{k=1}^{n} k^3 = 1^3 + 2^3 + 3^3 + \dots + n^3 = (1 + 2 + 3 + \dots + n)^2$$

c.
$$\sum_{k=1}^{n} k^3 = 1^3 + 2^3 + 3^3 + \dots + n^3 = (1 + 2 + 3 + \dots + n)^2$$

d. $\sum_{k=1}^{n} \frac{1}{(2k-1)(2k+1)} = \frac{1}{1\cdot 3} + \frac{1}{3\cdot 5} + \frac{1}{5\cdot 7} + \dots + \frac{1}{(2n-1)(2n+1)} < \frac{1}{2}$

e.
$$\sum_{k=1}^{n} \frac{1}{(7k-6)(7k+1)} = \frac{1}{1\cdot 8} + \frac{1}{8\cdot 15} + \frac{1}{15\cdot 22} + \dots + \frac{1}{(7n-6)(7n+1)} < \frac{1}{7}$$

f.
$$\sum_{k=n+1}^{3n+1} \frac{1}{k} = \frac{1}{n+1} + \frac{1}{n+2} + \frac{1}{n+3} + \dots + \frac{1}{3n+1} > 1$$

Recap. In order to prove the equality A(n) = B(n), for any n, using the method of mathematical induction you have to

- Prove that A(1) = B(1)
- Prove that A(k + 1) A(k) = B(k + 1) B(k)
- Then from assumption A(k) = B(k) and from equality (*) follows A(k + 1) = B(k + 1)
- Prove by mathematical induction that for any natural number n_i 3.
 - a. $5^n + 6^n 1$ is divisible by 10
 - b. $9^{n+1} 8n 9$ is divisible by 64
- **Recap.** Binomial coefficients are defined by 4.

$$C_n^k = {}_k C_n = {n \choose k} = \frac{n!}{k!(n-k)!}$$

- a. Prove that $C_{n+k}^2 + C_{n+k+1}^2$ is a full square
- b. Find *n* satisfying the following equation,

$$C_n^{n-1} + C_n^{n-2} + C_n^{n-3} + \dots + C_n^{n-10} = 1023$$

c. Prove that

$$\frac{C_n^1 + 2C_n^2 + 3C_n^3 + \dots + nC_n^n}{n} = 2^{n-1}$$

5. Find the roots of the equation:

$$\frac{14}{20-6x-2x^2} + \frac{x^2+4x}{x^2+5x} - \frac{x+3}{x-2} + 3 = 0.$$

Geometry.

Review the classwork handout. Solve the remaining problems from the previous homework; consider solutions explained in the classwork handout. Try solving the following additional problems. In all the problems, you are only allowed to use theorems we had proven before.

Problems.

- 1. Prove that for any triangle *ABC* with sides a, b and c, the area, $S \le \frac{1}{4}(b^2 + c^2)$.
- 2. In an isosceles triangle ABC with the side |AB| = |BC| = b, the segment |A'B'| = m connects the intersection points of the bisectors, AA' and BB' of the angles at the base, AC, with the corresponding opposite sides, $A' \in BC$ and $B' \in AB$. Find the length of the base, |AC| (express through given lengths, b and m).
- 3. Prove that for any point on a side of an equilateral triangle, the sum of the distances to the two other sides is the same constant. What is this distance (the side of the triangle is a)?
- 4. Distances from the point M inside an equilateral triangle ABC to the respective sides of this triangle are, d_a , d_b and d_c . Find the altitude of this triangle.
- 5. Three lines parallel to the respective sides of the triangle ABC intersect at a single point, which lies inside this triangle. These lines split the triangle ABC into 6 parts, three of which are triangles with areas S_1 , S_2 , and S_3 . Show that the area of the triangle ABC,

$$S = \left(\sqrt{S_1} + \sqrt{S_2} + \sqrt{S_3}\right)^2.$$