

## Algebra.

### Trigonometric form of complex numbers. Geometric interpretation.

Let us consider complex numbers with the absolute value of 1,

$$z_1 = x_1 + iy_1, |z_1|^2 = z_1 \bar{z}_1 = x_1^2 + y_1^2 = 1.$$

There is an obvious one-to-one correspondence between such numbers and points  $Z_1(x_1, y_1)$  on a circle of unit radius. Hence, we can express such numbers in terms of an angle,  $\varphi$ , parameterizing points on the unit circle,

$$z_1 = x_1 + iy_1 = \cos \varphi + i \sin \varphi.$$

More generally, any complex number,  $z = x + iy$ , whose absolute value is  $|z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2} = r$ , can be written in the trigonometric form as,

$$z = x + iy = r(\cos \varphi + i \sin \varphi).$$

Geometrically, it is represented by a point  $Z(x, y)$  on a circle of radius  $r = |z|$ . Position of this point is specified by an angle,  $\varphi$ , which is conventionally measured counterclockwise from the positive direction of the  $X$ -axis. Angle  $\varphi$  is called the argument of the complex number  $z$  and is denoted  $\varphi = \text{Arg}(z)$ . Thus, instead of describing a complex number by its real and imaginary part, i.e. its coordinates,  $(x, y)$ , we can describe it by its magnitude and argument (polar coordinates),  $(r, \varphi)$ , where  $r \geq 0$  and  $0 \leq \varphi = \text{Arg}(z) < 360^\circ$ .


It is now easy to prove the following important property of the multiplication of complex numbers.

**Theorem.** When we multiply two complex numbers, magnitudes multiply and arguments add,

$$|z_1 z_2| = |z_1| |z_2|, \text{Arg}(z_1 z_2) = (\text{Arg}(z_1) + \text{Arg}(z_2)) \text{mod } 360^\circ.$$

**Proof.** Let  $\text{Arg}(z_1) = \varphi_1$  and  $\text{Arg}(z_2) = \varphi_2$ , so  $z_1 = |z_1|(\cos \varphi_1 + i \sin \varphi_1)$  and  $z_2 = |z_2|(\cos \varphi_2 + i \sin \varphi_2)$ . Perform the multiplication directly,

$$z_1 z_2 = |z_1|(\cos \varphi_1 + i \sin \varphi_1) |z_2|(\cos \varphi_2 + i \sin \varphi_2) =$$



$$|z_1||z_2|(\cos \varphi_1 \cos \varphi_2 - \sin \varphi_1 \sin \varphi_2 + i(\sin \varphi_1 \cos \varphi_2 + \cos \varphi_1 \sin \varphi_2)) \\ = |z_1||z_2|(\cos(\varphi_1 + \varphi_2) + i \sin(\varphi_1 + \varphi_2))$$

Complex numbers whose arguments would differ by multiples of  $360^\circ$  are identical and correspond to the same point on the complex plane. Hence, the argument is computed *mod*  $360^\circ$ , ensuring that  $0 \leq \varphi_1 + \varphi_2 < 360^\circ$ .

**Theorem.** Multiplication of a complex number,  $z = x + iy = r(\cos \varphi + i \sin \varphi)$ , by a complex number of unit magnitude and argument  $\psi$ ,

$$z_\psi = x_\psi + iy_\psi = \cos \psi + i \sin \psi,$$

corresponds to a counterclockwise rotation of the point,  $Z(x, y)$ , on the complex plane, by an angle  $\psi$ ,

$$|zz_\psi| = r, \text{ Arg}(zz_\psi) = \varphi + \psi.$$

**Proof.** Indeed, perform the multiplication directly,

$$zz_\psi = (x + iy)(x_\psi + iy_\psi) = r(\cos \varphi + i \sin \varphi)(\cos \psi + i \sin \psi) \\ = r(\cos \varphi \cos \psi - \sin \varphi \sin \psi + i(\sin \varphi \cos \psi + \cos \varphi \sin \psi)) \\ = r(\cos(\varphi + \psi) + i \sin(\varphi + \psi))$$

It is clear that multiplication by a complex number with magnitude  $r'$  and argument  $\psi$  is equivalent to the combination of multiplication by a number of unit magnitude and argument  $\psi$ , and by the real number  $r'$ .

**Theorem.** Multiplication of a complex number,  $z = x + iy$ , by a complex number with magnitude  $r'$  and argument  $\psi$ ,

$$w = r'(\cos \psi + i \sin \psi),$$

results in a point on the complex plane, which is obtained from the point  $Z(x, y)$  by the combination of a rotation by angle  $\psi$  and a homothety (rescaling) with scale factor  $r'$ .

Multiplication of all complex numbers by a complex number  $w = r'(\cos \psi + i \sin \psi)$  is a transformation of the complex plane, which maps complex plane on itself. Identifying multiplication by a complex number with such transformation, we can state the following.

**Theorem.** Multiplication by a complex number with magnitude  $r'$  and argument  $\psi$ ,  $w = r'(\cos \psi + i \sin \psi)$ , is a combination of rotation by angle  $\psi$  and homothety (rescaling) with scale factor  $r'$ .

### De Moivre's formula.

**Theorem.** The formula named after Abraham de Moivre states that for any complex number,  $z = x + iy = r(\cos \varphi + i \sin \varphi)$ , and for any integer  $n \in \mathbb{N}$ ,

$$z^n = (r(\cos \varphi + i \sin \varphi))^n = r^n(\cos n\varphi + i \sin n\varphi)$$

**Proof 1 (Mathematical induction).**

1. Base case,  $n = 1$ :  $z^1 = r(\cos \varphi + i \sin \varphi)$  is true.
2.  $I(n) \Rightarrow I(n + 1)$ . Assume  $z^n = r^n(\cos n\varphi + i \sin n\varphi)$  is true. Then,

$$\begin{aligned} z^{n+1} &= z \cdot z^n = r(\cos \varphi + i \sin \varphi) \cdot (r(\cos \varphi + i \sin \varphi))^n \\ &= r(\cos \varphi + i \sin \varphi)r^n(\cos n\varphi + i \sin n\varphi) \\ &= r^{n+1}(\cos \varphi \cos n\varphi - \sin \varphi \sin n\varphi \\ &\quad + i(\sin \varphi \cos n\varphi + \cos \varphi \sin n\varphi)) \\ &= r^{n+1}(\cos(n + 1)\varphi + i \sin(n + 1)\varphi) \end{aligned}$$

**Proof 2 (Geometrical).**

$$z^n = (r(\cos \varphi + i \sin \varphi))^n = r(\cos \varphi + i \sin \varphi)r(\cos \varphi + i \sin \varphi) \dots r(\cos \varphi + i \sin \varphi)$$

By property of the multiplication of complex numbers, absolute values multiply, while arguments add. Therefore,

$$|z^n| = |z|^n = r^n, \text{ and } Arg(z^n) = nArg(z) = n\varphi, \text{ wherefrom it follows that } z^n = r^n(\cos n\varphi + i \sin n\varphi).$$

### n-th root.

The formula of de Moivre allows us to compute  $n$ -th root of a complex number. Suppose we want to solve the equation,

$$w^n = z$$

where  $w, z \in \mathbb{C}$ , so  $w$  is the  $n$ -th root of  $z$ . According to de Moivre's formula, if  $w = |w|(\cos \psi + i \sin \psi)$ , then  $w^n = |w|^n(\cos n\psi + i \sin n\psi)$ . Denoting  $z = r(\cos \varphi + i \sin \varphi)$ , we can rewrite the equation as,

$$w^n = |w|^n(\cos n\psi + i \sin n\psi) = r(\cos \varphi + i \sin \varphi)$$

One obvious solution is  $r = |w|^n$  and  $\varphi = n\psi$ ,  $w = \sqrt[n]{r} \left( \cos \frac{\varphi}{n} + i \sin \frac{\varphi}{n} \right)$ .

However, because  $\varphi = \text{Arg}(z) = \text{Arg}(w^n)$  and  $\psi = \text{Arg}(w)$  are determined modulo  $360^\circ$  ( $2\pi$  radians), there are other solutions, too, satisfying the above equation, such as  $w = \sqrt[n]{r} \left( \cos \frac{\varphi + 2\pi}{n} + i \sin \frac{\varphi + 2\pi}{n} \right)$ . Generally, we must have  $r = |w|^n$ , and  $\varphi = \text{Arg}(z) = \text{Arg}(w^n) \bmod 360^\circ = n\text{Arg}(w) \bmod 360^\circ = n\psi \bmod 360^\circ$ . Altogether, there are  $n$  solutions,

$$w = \sqrt[n]{r} \left( \cos \frac{\varphi + 2\pi k}{n} + i \sin \frac{\varphi + 2\pi k}{n} \right), 0 \leq k < n$$

This is a special case of the following extremely important result, called the fundamental theorem of algebra.

**Theorem.** Any polynomial with complex coefficients of degree  $n$  has exactly  $n$  roots (counting with multiplicities).

There is no simple proof of this theorem (and, in fact, no purely algebraic proof: all the known proofs use some geometric arguments).

In particular, since any polynomial with real coefficients can be considered as a special case of a polynomial with complex coefficients, this shows that any real polynomial of degree  $n$  has exactly  $n$  complex roots.