

**MATH 7**  
**HOMEWORK 11: VIETA FORMULAS AND QUADRATIC INEQUALITIES**  
DEC 8, 2018

VIETA FORMULAS

If a polynomial  $p(x)$  has root  $a$  (i.e., if  $p(a) = 0$ ), then  $p(x)$  is divisible by  $(x - a)$ , i.e.  $p(x) = (x - a)q(x)$  for some polynomial  $q(x)$ . In particular, if  $x_1, x_2$  are roots of quadratic polynomial  $ax^2 + bx + c$ , then  $ax^2 + bx + c = a(x - x_1)(x - x_2)$ .

Therefore, if  $a = 1$ , then

$$\begin{aligned}x_1 + x_2 &= -b \\ x_1 x_2 &= c\end{aligned}$$

(Vieta formulas).

SOLVING POLYNOMIAL INEQUALITIES

We discussed the general rule for solving polynomial inequalities:

- Find the roots and factor your polynomial, writing it in the form  $p(x) = a(x - x_1)(x - x_2)$  (for polynomial of degree more than 2, you would have more factors).
- Roots  $x_1, x_2, \dots$  divide the real line into intervals; define the sign of each factor and the product on each of the intervals.

HOMEWORK

1. Can you guess an analog of Vieta formulas for equation of degree 3: if  $x_1, x_2, x_3$  are roots of an equation  $x^3 + bx^2 + cx + d$ , then what is the relation between  $b, c, d$  and  $x_1, x_2, x_3$ ?
2. Let  $x_1, x_2$  be roots of equation  $x^2 + 5x - 7 = 0$ . Find  
(a)  $x_1^2 + x_2^2$  (b)  $(x_1 - x_2)^2$  (c)  $\frac{1}{x_1} + \frac{1}{x_2}$  (d)  $x_1^3 + x_2^3$   
(hint for part (d): compute first  $(x_1 + x_2)(x_1^2 + x_2^2)$  )
- \*3. Prove the statement we used in class: if a polynomial  $p(x)$  has root  $a$  (i.e., if  $p(a) = 0$ ), then  $p(x)$  is divisible by  $(x - a)$ , i.e.  $p(x) = (x - a)q(x)$  for some polynomial  $q(x)$ .
4. Solve the equation  $x^4 - 3x^2 + 2 = 0$ .
5. Solve the following equations and inequalities:

$$\begin{array}{lll} \text{(a) } x^2 - 5x + 6 > 0 & \text{(b) } x^2 < 1 + x & \text{(c) } \frac{x+1}{x-2} > 0 \\ \text{(d) } x(x-5)(x+7) < 0 & \text{(e) } \sqrt{2x+1} = x & \text{(f) } \frac{2x+1}{x-3} > 1 \end{array}$$

6. (a) Show that for any  $a, b \geq 0$ , one has  $\frac{a+b}{2} \geq \sqrt{ab}$ . (The left hand side is usually called the arithmetic mean of  $a, b$ ; the right hand side is called the geometric mean of  $a, b$ .)  
(b) Prove that for any  $a > 0$ , we have  $a + \frac{1}{a} \geq 2$ , with equality only when  $a = 1$ .
- \*7. Solve equation  $x^4 - 5x^3 + 6x^2 - 5x + 1 = 0$ . [Hint: divide by  $x^2$  and try to rewrite as an equation in  $y = x + (1/x)$ . The same trick works for any symmetric equation, in which coefficient of  $x^4$  is the same as the constant term, and coefficient of  $x^3$  is the same as coef. of  $x$ . ]
8. For what values of  $a$  does  $x^2 + ax + 14$  has no roots? exactly one root? two roots?