## MATH 7 HOMEWORK 13: PARABOLA JAN 6, 2018

GRAPHS OF y = |x| and  $y = x^2$ 

The figure below shows graphs of functions y = |x| and  $y = x^2$ ; the latter graph is called a *parabola*.



TRANSFORMATIONS OF GRAPHS

- Graph of function y = f(x + a) is obtained from graph of y = f(x) by shifting horizontally by a units to the left; for example, graph of  $y = (x+1)^2$  is parabola with vertex at (-1, 0).
- Graph of function y = f(x) + k is obtained from graph of y = f(x) by shifting vertically by k units up; for example, graph of  $y = x^2 + 1$  is parabola with vertex at (0, 1).
- Finally, graph of y = kf(x) is obtained from graph of y = f(x) by rescaling vertically by factor of k; if k is negative, it means flip upside down and then rescale by factor of |k|.

Combining these results, we can sketch the graph of any quadratic function, which will also be a parabola. To sketch it, we need to complete the square, writing

$$ax^{2} + bx + c = a\left(x + \frac{b}{2a}\right)^{2} - \frac{b^{2} - 4ac}{4a} = a(x - h)^{2} + k, \qquad h = -\frac{b}{2a}, \quad k = -\frac{b^{2} - 4ac}{4a}$$

For example:  $x^2 + x = (x + \frac{1}{2})^2 - \frac{1}{4}$ 

The result will be a parabola obtained by stretching the usual parabola vertically by factor a (if a < 0, this means flipping it upside down and then stretching by |a|) and then moving it so that the vertex will be at point (h, k),

In particular, the branches go up if a > 0 and down if a < 0.

## Homework

- 1. For what values of a does the polynomial  $x^2 + ax + 14$  has no roots? exactly one root? two roots?
- 2. Sketch the graphs of the following functions and relations:
  - (a) x + y = 4(b) |x| + y = 4(c)  $x^2 + 4x + y^2 4y = 0$ (d) y = |x 5|(e) y = |x + 1| + |x 1|(f)  $y = x^2 x$ (g)  $y = |x^2 x|$ (h)  $y = x^2 5x + 6$ (i)  $y = -2x^2 + 8x + 6$
- 3. Solve the following equations and inequalities
  - (a)  $x^2 x + 6 \ge 0$  (b)  $\frac{2x+1}{x-5} \le 0$  (c)  $x^4 3x^2 + 8 = 0$ (d) x(x-2)(x+18) > 0
- 4. Find all intersection points of parabola  $y = x^2$  and the circle with radius  $\sqrt{6}$  and center at (0, 4).
- 5. Prove that for any point P on the parabola  $y = \frac{x^2}{4} + 1$ , the distance from P to the x-axis is equal to the distance from P to the point (0, 2).