

Homework 27: Trigonometry: radians and the unit circle.


HW27 is Due May 10; submit it to Google Classroom 15 minutes before the class time.

1. Definition for sin, cos, and tan of an angle

In general, for a right-angle triangle, the **sin** α and **cos** α of the angle are defined as:

$$\sin(\alpha) = \frac{\text{opposite side}}{\text{hypotenuse}} = \frac{a}{c}, \quad \cos(\alpha) = \frac{\text{adjacent side}}{\text{hypotenuse}} = \frac{b}{c}$$

$$\tan(\alpha) = \frac{\sin(\alpha)}{\cos(\alpha)} = \frac{\text{opposite side/hypotenuse}}{\text{adjacent side/hypotenuse}} = \frac{a}{b}$$



2. Radians – a new measure of the angle's size

Until now, we have been measuring angles in degrees, which are defined by saying that a full turn corresponds to 360° . An alternative way to measure angles is by **radians**, which are defined in the following way: given an angle θ , its measure in radians is the ratio of the arc (**a**) of the circumference and the radius (**r**).

For example, the angle 360° corresponds to a full circle. Since the perimeter of a circle is $2\pi R$, dividing by R gives: **$360^\circ = 2\pi \text{ rad}$** . In the same way, half a circle corresponds to an angle of $180^\circ = \frac{1/2 \times 2\pi R}{R} = \pi$ radians. By similar arguments, we can translate all the angles that appeared in our previous table:




FIGURE 2.

| Function | Notation | Definition | 0° | 30° | 45° | 60° | 90° |
|----------|----------------|---|-----------|----------------------|----------------------|----------------------|------------|
| | | angle in rad | 0 | $\pi/6$ | $\pi/4$ | $\pi/3$ | $\pi/2$ |
| sine | $\sin(\alpha)$ | $\frac{\text{opposite side}}{\text{hypotenuse}}$ | 0 | $\frac{1}{2}$ | $\frac{\sqrt{2}}{2}$ | $\frac{\sqrt{3}}{2}$ | 1 |
| cosine | $\cos(\alpha)$ | $\frac{\text{adjacent side}}{\text{hypotenuse}}$ | 1 | $\frac{\sqrt{3}}{2}$ | $\frac{\sqrt{2}}{2}$ | $\frac{1}{2}$ | 0 |
| tan | $\tan(\alpha)$ | $\frac{\text{opposite side}}{\text{adjacent side}}$ | 0 | $\frac{1}{\sqrt{3}}$ | 1 | $\sqrt{3}$ | ∞ |

3. Trigonometric (unit) circle

A very useful tool for understanding trigonometric functions is the trigonometric circle (see Figure 3). To find the sine and cosine of a positive angle α , we just have to “walk” around the circle a distance α , starting from the point (1; 0) in an anticlockwise direction. Then, the coordinates of the point we arrive at are $(\cos \alpha; \sin \alpha)$. For negative angles α , we define the sine and cosine in the same way, but walking in a clockwise direction.

Note, that you could have angles larger than 360° . For example, the angle $390^\circ = 360^\circ + 30^\circ$ that has the same coordinates on the unit circle as a 30° angle (you must circle counterclockwise 1 full revolution and 30 more degrees).




FIGURE 3.

Homework problems

Instructions: Please always write solutions on a **separate sheet of paper**. Solutions should include explanations. I want to see more than just an answer: I also want to see how you arrived at this answer, and some justification why this is indeed the answer. So **please include sufficient explanations**, which should be clearly written so that I can read them and follow your arguments.

All angles are measured in degrees or radians.

1. Draw a large unit circle or use the attached. Then, remembering that 2π rad corresponds to a full circle, find the points on the circumference of the circle that correspond to the following angles. Label the angle with the *rad* value.

- | | | |
|---------------------|----------------------|-----------------------|
| a) π | d) $-\frac{5\pi}{4}$ | g) $\frac{25\pi}{3}$ |
| b) $\frac{3\pi}{2}$ | e) 11π | h) $-\frac{19\pi}{6}$ |
| c) $\frac{3\pi}{4}$ | f) -3π | |

2. Using a calculator, compare the results. Be careful with the setting (DEG or RAD) of your calculator. If the angle is measured in degrees, use DEG. If the angle is measured in radians, use RAD.

- | | | | | |
|---------------------|-----|-----------------------|-----|----------------------|
| a. $\sin 45^\circ$ | and | c. $\sin 90^\circ$ | and | d. $\cos 0^\circ$ |
| b. $\cos 120^\circ$ | and | $\sin \frac{\pi}{4}$ | and | $\sin \frac{\pi}{2}$ |
| | | $\cos \frac{2\pi}{3}$ | | $\cos 0$ |

3. Now, use your unit circle, Figure 3, and the values for sin/cos from the table on page 1 to complete this table:

| Angle in rad | sine | cosine |
|--------------|------|--------|
| π | | |
| $3\pi/2$ | | |
| $3\pi/4$ | | |
| $-5\pi/2$ | | |
| 11π | | |
| -3π | | |
| $25\pi/3$ | | |
| $-19\pi/6$ | | |


4. (* Optional) Using the unit circle, check the interval where the inequalities are valid for the given angles x .

| angle x | $\sin x \geq \frac{\sqrt{3}}{2}$ | $\frac{1}{2} < \sin x < \frac{\sqrt{3}}{2}$ | $-\frac{\sqrt{2}}{2} < \sin x < \frac{1}{2}$ | $\sin x \leq -\frac{\sqrt{2}}{2}$ |
|-----------|----------------------------------|---|--|-----------------------------------|
| $\pi/7$ | | | ✓ | |
| $2\pi/7$ | | | | |
| $-3\pi/5$ | | | | |
| $5\pi/8$ | | | | |
| $25\pi/9$ | | | | |

5. Using the trigonometric circle, show that $\cos x = \sin(x + \pi/2)$ for any angle x .

6. (* Optional) Find all real numbers x such that $(\sin x)^2 = \frac{3}{4}$ in the interval 0 to 2π . Are there more angles like that outside this interval?

You may find this unit circle helpful for problem 1, or you may draw your own.



This circle may help you with problems 3 and 4. The values in brackets are the (x,y) coordinates of the point where your angle ends (terminal point). They will represent the values of the $(\cos\alpha, \sin\alpha)$. You could fill in this circle on your own by using the values from the table on page 1 or the unit circle from *desmos* which we used in class (look in the Class Notes folder).

