


MATH 8
HANDOUT 6: LOGIC 1: INTRODUCTION TO SYMBOLS AND FORMULAS



Today we will start discussing formal rules of logic.

In logic, we will be dealing with *boolean* expressions, i.e. expressions which only take two values, TRUE and FALSE. We will commonly use abbreviations T and F for these values.

You can also think of these two values as the two possible digits in binary (base 2) arithmetic: $T = 1$, $F = 0$.

In the usual arithmetic, we have some operations (addition, multiplication, ...) which satisfy certain laws (associativity, distributivity, ...). Similarly, there are logic operations and logic laws.

BASIC LOGIC OPERATIONS

NOT (for example, NOT A): true if A is false, and false if A is true. Commonly denoted by $\neg A$ or (in computer science) $!A$.

AND (for example A AND B): true if both A, B are true, and false otherwise (i.e., if at least one of them is false). Commonly denoted by $A \wedge B$

OR (for example A OR B): true if at least one of A, B is true, and false otherwise. Sometimes also called “inclusive or” to distinguish it from the “exclusive or” described in problem 4 below. Commonly denoted by $A \vee B$.

IF (as in “if A , then B ”; written $A \implies B$): if A is false, automatically true; if A is true, it is true only when B is true

As in usual algebra, logic operations can be combined, e.g. $(A \vee B) \wedge C$.

TRUTH TABLES

If we have a logical formula involving variables A, B, C, \dots , we can make a table listing, for every possible combination of values of A, B, \dots , the value of our formula. For example, the following is the truth tables for OR and IF:

A	B	A OR B
T	T	T
T	F	T
F	T	T
F	F	F

A	B	A AND B
T	T	T
T	F	F
F	T	F
F	F	F

LOGIC LAWS

We can combine logic operations, creating more complicated expressions such as $A \wedge (B \vee C)$. As in arithmetic, these operations satisfy some laws: for example $A \vee B$ is the same as $B \vee A$. Here, “the same” means “for all values of A, B , these two expressions give the same answer”; it is usually denoted by \iff . Here are two other laws:

$$\begin{aligned}\neg(A \wedge B) &\iff (\neg A) \vee (\neg B) \\ A \wedge (B \vee C) &\iff (A \wedge B) \vee (A \wedge C)\end{aligned}$$

Truth tables provide the most straightforward (but not the shortest) way to prove complicated logical rules: if we want to prove that two formulas are equivalent (i.e., always give the same answer), make a truth table for each of them, and if the tables coincide, they are equivalent.

PROBLEMS

- Write the truth table for each of the following formulas. Are they equivalent (i.e., do they always give the same value)?
 - $(A \vee B) \wedge (A \vee C)$
 - $A \vee (B \wedge C)$.

- Use the truth tables to prove *De Morgan’s laws*

$$\begin{aligned}\neg(A \wedge B) &\iff (\neg A) \vee (\neg B) \\ \neg(A \vee B) &\iff (\neg A) \wedge (\neg B)\end{aligned}$$

- Use truth tables to show that \vee is commutative and associative:

$$\begin{aligned}A \vee B &\iff B \vee A \\ A \vee (B \vee C) &\iff (A \vee B) \vee C\end{aligned}$$

Is it true that \wedge is also commutative and associative?

- Another logic operation, called “exclusive or”, or XOR, is defined as follows: $A \text{ XOR } B$ is true if and only if exactly one of A, B is true.
 - Write a truth table for XOR
 - Describe XOR using only basic logic operations AND, OR, NOT, i.e. write a formula using variable A, B and these basic operations which is equivalent to $A \text{ XOR } B$.
- Yet one more logic operation, NAND, is defined by

$$A \text{ NAND } B \iff \text{NOT}(A \text{ AND } B)$$

- Write a truth table for NAND
- What is $A \text{ NAND } A$?

*

- Show that you can write NOT A , $A \text{ AND } B$, $A \text{ OR } B$ using only NAND (possibly using each of A, B more than once).

This last part explains why NAND chips are popular in electronics: using them, you can build **any** logical gates.

- A restaurant menu says *The fixed price dinner includes entree, dessert, and soup or salad.*

Can you write it as a logical statement, using the following basic pieces:

E : your dinner includes an entree

D : your dinner includes a dessert

P : your dinner includes a soup

S : your dinner includes a salad

and basic logic operations described above?

- On the island of knights and knaves, there are two kinds of people: Knights, who always tell the truth, and Knaves, who always lie. Unfortunately, there is no easy way of knowing whether a person you meet is a knight or a knave...

You meet two people on this island, Bart and Ted. Bart claims, “I and Ted are both knights or both knaves.” Ted tells you, “Bart would tell you that I am a knave.” So who is a knight and who is a knave?