Math 6b/c: Homework 24 Homework #24 is due April 29.

## Geometric progression

A sequence of numbers is a geometric progression if the next number in the sequence is the current number times a constant called the common ratio, let's call it q. For example, let's consider the sequence:

The first term in the sequence is  $b_1 = 6$ , the second is  $b_2 = 6 \times 2 = 12$ , and so on.

The common ratio is q = 2. Indeed  $b_3 = b2 \times q = 12 \times 2 = 24$  and  $b_4 = 24 \times 2 = 48$ .

What is the  $n^{th}$  term? For example what is  $b_{10}$ ?

$$b_{1} = 6$$
  

$$b_{2} = b_{1} \times q = 6 \times 2 = 12$$
  

$$b_{3} = b_{2} \times q = (b_{1} \times q) \times q = b_{1} \times q^{2} = 6 \times 2^{2} = 24$$
  

$$b_{4} = b_{3} \times q = (b_{1} \times q^{2}) \times q^{2} = b_{1} \times q^{3} = 6 \times 2^{3} = 48$$
  
....  

$$b_{n} = b_{1} \times q^{n-1}$$
  
So  $b_{10} = b_{1} \times q^{9} = 6 \times 2^{9} = 6 \times 512 = 3072$ 

## Sum of a geometric progression

There is a formula for the sum of a geometric progression:

$$S = b_1 + b_2 + b_3 + \dots + b_n = b_1 \times \frac{(1 - q^n)}{1 - q}$$

To prove this, we write the sum and we multiply it by q:

$$S = b_1 + b_2 + b_3 + \dots + b_{n-1} + b_n$$
$$qS = qb_1 + qb_2 + qb_3 + \dots + qb_{n-1} + qb_n$$

Remember that  $qb_{n-1} = b_n$ , so that the last term is  $qb_n = q \times (b_1 \times q^{n-1}) = b_1 \times q^n$ .

$$qS = b_2 + b_3 + b_4 + \dots + b_n + b_1 q^n$$

We subtract *S* from each side:

$$qS - S = b_2 + b_3 + \dots + b_n + b_1 q^n - (b_1 + b_2 + b_3 + \dots + b_{n-1} + b_n)$$

All terms cancel, except  $b_1q^n$  and  $b_1$  so that:

$$qS - S = b_1 q^n - b_1$$
$$(1 - q)S = b_1 q^n - b_1$$
$$S = \frac{b_1 q^n - b_1}{1 - q} = \frac{b_1 (1 - q^n)}{1 - q}$$

## **Homework**

- 1. Write the first 5 terms of a geometric progression if  $b_1 = -20$  and  $q = \frac{1}{2}$
- 2. What are the first 2 terms of the geometric progression:  $b_1$ ,  $b_2$ , 24, 36, 54, ...?
- 3. What is the common ratio of the geometric progression:  $\frac{1}{2}$ ,  $-\frac{1}{2}$ ,  $\frac{1}{2}$ ,  $-\frac{1}{2}$ , ...? What is  $b_{10}$ ? What is  $b_{100}$ ?
- 4. Simplify:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^{10}}$$

- 5. What is the sum:  $1 2 + 2^2 2^3 + 2^4 2^5 + \dots 2^{15}$ ?
- 6. What is the sum:  $1 + x + x^2 + x^3 + x^4 + x^5 + \dots + x^{100}$ ?
- 7. A geometric progression has 99 terms, the first term is 12 and the last term is 48. What is the 50th term?
- 8. If we put one grain of wheat on the first square of the chessboard, two on the second, then four, eight, . . , approximately how many grains of wheat will there be? (You can use  $2^{10} = 1024 \approx 10^3$ ). Can you estimate the total volume of all this wheat? Compare with the annual wheat harvest of the US, which is about 2 billion bushels. (A grain of wheat is about 10 mm<sup>3</sup>; a bushel is about 35 liters, or 0.035 m<sup>3</sup>)
- 9. How many multiples of 7 are there between 1 and 1000? Can you find the sum of them all?
- 10. Find the sum  $1 + 3 + 5 + \cdots + 999$ .