

MATH 7
VIETA FORMULAS AND QUADRATIC INEQUALITIES

VIETA FORMULAS

If a polynomial $p(x)$ has root a (i.e., if $p(a) = 0$), then $p(x)$ is divisible by $(x - a)$, i.e. $p(x) = (x - a)q(x)$ for some polynomial $q(x)$. In particular, if x_1, x_2 are roots of quadratic polynomial $ax^2 + bx + c$, then $ax^2 + bx + c = a(x - x_1)(x - x_2)$.

Therefore, if $a = 1$, then

$$\begin{aligned}x_1 + x_2 &= -b \\x_1x_2 &= c\end{aligned}$$

(Vieta formulas).

SOLVING POLYNOMIAL INEQUALITIES

We discussed the general rule for solving polynomial inequalities:

- Find the roots and factor your polynomial, writing it in the form $p(x) = a(x - x_1)(x - x_2)$ (for polynomial of degree more than 2, you would have more factors).
- Roots x_1, x_2, \dots divide the real line into intervals; define the sign of each factor and the product on each of the intervals.

1. Can you guess an analog of Vieta formulas for equation of degree 3: if x_1, x_2, x_3 are roots of an equation $x^3 + bx^2 + cx + d$, then what is the relation between b, c, d and x_1, x_2, x_3 ?
2. Let x_1, x_2 be roots of equation $x^2 + 5x - 7 = 0$. Find
(a) $x_1^2 + x_2^2$ (b) $(x_1 - x_2)^2$ (c) $\frac{1}{x_1} + \frac{1}{x_2}$ (d) $x_1^3 + x_2^3$
(hint for part (d): compute first $(x_1 + x_2)(x_1^2 + x_2^2)$)
- *3. Prove the statement we used in class: if a polynomial $p(x)$ has root a (i.e., if $p(a) = 0$), then $p(x)$ is divisible by $(x - a)$, i.e. $p(x) = (x - a)q(x)$ for some polynomial $q(x)$.
4. Solve the equation $x^4 - 3x^2 + 2 = 0$.
5. Solve the following equations and inequalities:

$$\begin{array}{lll} \text{(a) } x^2 - 5x + 6 > 0 & \text{(b) } x^2 < 1 + x & \text{(c) } \frac{x+1}{x-2} > 0 \\ \text{(d) } x(x-5)(x+7) < 0 & \text{(e) } \sqrt{2x+1} = x & \text{(f) } \frac{2x+1}{x-3} > 1 \end{array}$$

6. (a) Show that for any $a, b \geq 0$, one has $\frac{a+b}{2} \geq \sqrt{ab}$. (The left hand side is usually called the arithmetic mean of a, b ; the right hand side is called the geometric mean of a, b .)
(b) Prove that for any $a > 0$, we have $a + \frac{1}{a} \geq 2$, with equality only when $a = 1$.
- *7. Solve equation $x^4 - 5x^3 + 6x^2 - 5x + 1 = 0$. [Hint: divide by x^2 and try to rewrite as an equation in $y = x + (1/x)$. The same trick works for any symmetric equation, in which coefficient of x^4 is the same as the constant term, and coefficient of x^3 is the same as coef. of x .]
8. For what values of a does $x^2 + ax + 14$ has no roots? exactly one root? two roots?