

# MATH 8: ASSIGNMENT 15

JANUARY 20, 2018

## 1. PERPENDICULAR BISECTOR

Consider any property of points on the plane — for example, the property that a point  $P$  is a distance exactly  $r$  from a given point  $O$ . The set of all points  $P$  for which this property holds true is called the **locus** of points satisfying this property. As we have seen above, the locus of points that are a distance  $r$  from a point  $O$  is called a circle (specifically, a circle of radius  $r$  centered at  $O$ ).


Now consider we are given two points  $A, B$ . If a point  $P$  is an equal distance from  $A, B$  (i.e., if  $\overline{PA} \cong \overline{PB}$ ) then we say  $P$  is **equidistant** from points  $A, B$ .

**Theorem 14.** *The locus of points equidistant from a pair of points  $A, B$  is a line  $l$  which is perpendicular to  $\overline{AB}$  and goes through the midpoint of  $AB$ . This line is called the **perpendicular bisector** of  $\overline{AB}$ .*

*Proof.* Let  $M$  be the midpoint of  $\overline{AB}$ , and let  $l$  be the line through  $M$  which is perpendicular to  $AB$ . We need to prove that for any point  $P$ ,

$$(AP \cong BP) \iff P \in l$$

1. Assume that  $AP \cong BP$ . Then triangle  $APB$  is isosceles; by Theorem 10 from last week, it implies that  $PM \perp AB$ . Thus,  $PM$  must coincide with  $l$ , i.e.  $P \in l$ . Therefore, we have proved implication one way: if  $AP \cong BP$ , then  $P \in l$ .
2. Conversely, assume  $P \in l$ . Then  $m\angle AMP = m\angle BMP = 90^\circ$ ; thus, triangles  $\triangle AMP$  and  $\triangle BMP$  are congruent by SAS, and therefore  $AP \cong BP$ .



□

**Theorem 15.** *In a triangle  $\triangle ABC$ , the perpendicular bisectors of the 3 sides intersect at a single point. This point is the center of a circle circumscribed about the triangle (i.e., such that all three vertices of the triangle are on the circle).*

## 2. MEDIAN, ALTITUDE, ANGLE BISECTOR

Last week we defined three special lines that can be constructed from any vertex in any triangle; each line goes from a vertex of the triangle to the line containing the triangle's opposite side (altitudes may sometimes land on the opposite side outside of the triangle).


Given a triangle  $\triangle ABC$ ,

- The **altitude** from  $A$  is the line through  $A$  perpendicular to  $\overleftrightarrow{BC}$ ;
- The **median** from  $A$  is the line from  $A$  to the midpoint  $D$  of  $\overline{BC}$ ;
- The **angle bisector** from  $A$  is the line  $\overleftrightarrow{AE}$  such that  $\angle BAE \cong \angle CAE$ . Here we let  $E$  denote the intersection of the angle bisector with  $\overline{BC}$ .

The following result is an analog of theorem 14. For a point  $P$  and a line  $l$ , we define the distance from  $P$  to  $l$  to be the length of the perpendicular dropped from  $P$  to  $l$  (see problem 1 in the HW). We say that point  $P$  is equidistant from two lines  $l, m$  if the distance from  $P$  to  $l$  is equal to the distance from  $P$  to  $m$ .

**Theorem 16.** *For an angle  $ABC$ , the locus of points inside the angle which are equidistant from the two sides  $BA, BC$  is the ray  $\overrightarrow{BD}$  which is the angle bisector of  $\angle ABC$ .*

Proof of this theorem was discussed in class.




## 3. HOMEWORK

1. Let  $P$  be a point not on line  $l$ , and  $A \in l$  be the base of perpendicular from  $P$  to  $l$ :  $AP \perp l$ . Prove that for any other point  $B$  on  $l$ ,  $PB > PA$  ("perpendicular is the shortest distance"). Note: you can


not use Pythagorean theorem as we have not proved it yet; instead, try using Theorem 11 (opposite the larger angle there is a longer side).

2. Let  $\triangle ABC$  be a right triangle with right angle  $\angle A$ , and let  $D$  be the intersection of the line parallel to  $\overline{AB}$  through  $C$  with the line parallel to  $\overline{AC}$  through  $B$ .

- Prove  $\triangle ABC \cong \triangle DCB$
- Prove  $\triangle ABC \cong \triangle BDA$
- Prove that  $\overline{AD}$  is a median of  $\triangle ABC$ .



- Let  $\triangle ABC$  be a right triangle with right angle  $\angle A$ , and let  $D$  be the midpoint of  $\overline{BC}$ . Prove that  $AD = \frac{1}{2}BC$ .
- Let  $l_1, l_2$  be the perpendicular bisectors of side  $AB$  and  $BC$  respectively of  $\triangle ABC$ , and let  $F$  be the intersection point of  $l_1$  and  $l_2$ . Prove that then  $F$  also lies on the perpendicular bisector of the side  $BC$ . [Hint: use Theorem 14.]
- Prove Theorem 15.
- Let the angle bisectors from  $B$  and  $C$  in the triangle  $\triangle ABC$  intersect each other at point  $F$ . Prove that  $\overleftrightarrow{AF}$  is the third angle bisector of  $\triangle ABC$ . [Hint: use Theorem 16]
- Given triangle  $\triangle ABC$ , draw through each vertex a line parallel to the opposite side. Denote the vertices of the resulting triangle by  $D, E, F$ , as shown in the figure below.



- Prove that  $\triangle ABC \cong \triangle BAF$  (pay attention to order of vertices). Similarly one proves that all four small triangles in the picture are congruent.
- Prove that  $\overline{AB} \parallel \overline{ED}$  and  $AB = \frac{1}{2}ED$ .
- Prove that perpendicular bisectors of sides of  $\triangle DEF$  are altitudes of  $\triangle ABC$ .
- Show that in any triangle, the three altitudes meet at a single point.