

MATH 8: ASSIGNMENT 17

FEB 10, 2019

CIRCLES

Given a circle λ with center O ,

- A radius is any line segment from O to a point A on λ ,
- A chord is any line segment between distinct points A, B on λ ,
- A diameter is a chord that passes through O ,
- A tangent line is a line that intersects the circle exactly once; if the intersection point is A , the tangent is said to be the tangent through A .

Moreover, we say that two circles are tangent if they intersect at exactly one point.

Theorem 20. *Let A be a point on circle λ centered at O , and m a line through A . Then m is tangent to λ if and only if $m \perp \overline{OA}$. Moreover, there is exactly one tangent to λ at A .*

Proof. First we prove $(m \text{ is tangent to } \lambda) \implies (m \perp \overline{OA})$. Suppose m is tangent to λ at A but not perpendicular to \overline{OA} . Let \overline{OB} be the perpendicular to m through O , with B on m . Construct point C on m such that $BA = BC$; then we have that $\triangle OBA \cong \triangle OBC$ by *SAS*, using $OB = OB$, $\angle OBA = \angle OBC = 90^\circ$, and $BA = BC$. Therefore $OC = OA$ and hence C is on λ . But this means that m intersects λ at two points, which is a contradiction.

Now we prove $(m \perp \overline{OA}) \implies (m \text{ is tangent to } \lambda)$. Suppose m passes through A on λ such that $m \perp \overline{OA}$. If m also passed through B on λ , then $\triangle AOB$ would be an isosceles triangle since $\overline{AO}, \overline{BO}$ are radii of λ . Therefore $\angle ABO = \angle BAO = 90^\circ$, i.e. $\triangle AOB$ is a triangle with two right angles, which is a contradiction. \square

Notice that, given point O and line m , the perpendicular \overline{OA} from O to m (with A on m) is the shortest distance from O to m , therefore the locus of points of distance exactly OA from O should line entirely on one side of m . This is essentially the idea of the above proof.


Theorem 21. *Let \overline{AB} be a chord of circle λ with center O . Then O lies on the perpendicular bisector of \overline{AB} . Moreover, if C is on \overline{AB} , then C bisects \overline{AB} if and only if $\overline{OC} \perp \overline{AB}$.*

Proof. Let m be the perpendicular bisector of \overline{AB} . The center O of λ is equidistant from A, B by the definition of a circle, therefore by Theorem 14, O must be on m . Let m intersect \overline{AB} at D . We then have that D is the midpoint of \overline{AB} and also the foot of the perpendicular from O to \overline{AB} .

Then if C bisects \overline{AB} , C lies on the perpendicular bisector m of \overline{AB} , which passes through O , thus $\overline{OC} \perp \overline{AB}$. Lastly if $\overline{OC} \perp \overline{AB}$, then because there is only one perpendicular to \overline{AB} through O , we must have $C = D$ and hence C is the midpoint of \overline{AB} . \square

Theorem 22. *Let λ, ω be circles that intersect at points A, B . Then $\overline{AB} \perp \overline{OP}$.*

Proof. We have that $\triangle AOB$ and $\triangle APB$ are both isosceles, thus their altitudes from O and P respectively both intersect \overline{AB} at the midpoint C of \overline{AB} . Then, since $m\angle OCA = m\angle ACP = 90^\circ$, we have that $m\angle OCP = 180^\circ$, i.e. C lies on the line \overline{OP} . Since C is the foot of altitudes from O and P , this completes the proof. \square




Theorem 23. *Let λ, ω be circles that are both tangent to line m at point A . Then λ, ω are tangent circles.*

Proof. Suppose, by contradiction, that λ, ω intersect at point $B \neq A$. Then $\overline{AB} \perp \overline{OP}$, therefore both \overline{AB} and m are perpendicular to \overline{OA} through A . We must therefore have that B is on m , but m is tangent to λ through A , thus has only intersection with λ , which is a contradiction. \square

ARCS AND ANGLES

Consider a circle λ with center O , and an angle formed by two rays from O . Then these two rays intersect the circle at points A, B , and the portion of the circle contained inside this angle is called the **arc subtended** by $\angle AOB$.

Theorem 24. *Let A, B, C be on circle λ with center O . Then $\angle ACB = \frac{1}{2}\angle AOB$. The angle $\angle ACB$ is said to be inscribed in λ .*



Proof. There are actually a few cases to consider here, since C may be positioned such that O is inside, outside, or on the angle $\angle ACB$. We will prove the first case here, which is pictured on the left.

Case 1. Draw in segment \overline{OC} and notice that $m\angle AOC + m\angle BOC + m\angle AOB = 360^\circ$. Since \overline{OC} is a radius of λ , we have that $\triangle AOC$ and $\triangle BOC$ are isosceles triangles, thus $m\angle AOC = 180^\circ - 2m\angle OCA$ and $m\angle BOC = 180^\circ - 2m\angle OCB$. Therefore we get $180^\circ - 2m\angle ACO + 180^\circ - 2m\angle BCO + m\angle AOB = 360^\circ \implies m\angle AOB = 2(m\angle ACO + m\angle BCO) \implies m\angle AOB = 2m\angle ACB$. \square

As a result of Theorem 24, we get that any triangle $\triangle ABC$ on λ where \overline{AB} is a diameter must be a right triangle, since the angle $\angle ACB$ has half the measure of angle $\angle AOB$, which is 180° .

The idea captured by the concept of an arc and Theorem 24 is that there is a fundamental relationship between angles and arcs of circles, and that the angle 360° can be thought of as a full circle around a point.

HOMEWORK

1. Prove that, given a segment \overline{AB} , there is a unique circle with diameter \overline{AB} .
2. Given lines $\overleftrightarrow{AB} \parallel \overleftrightarrow{CD}$ such that $\overline{AD}, \overline{BC}$ intersect at E and $AE = ED$, prove that $BE = EC$.
3. Prove that if a diameter of circle λ is a radius of circle ω , then λ, ω are tangent.
4. Complete the proof of Theorem 24 by proving the cases where O is not inside the angle $\angle ACB$. [Hint: for one of the cases, you may need to write $\angle ACB$ as the difference of two angles.]
5. Prove the converse of Theorem 24: namely, if λ is a circle centered at O and A, B , are on λ , and there is a point C such that $m\angle ACB = \frac{1}{2}m\angle AOB$, then C lies on λ . [Hint: we need to prove that $OC = OA$; consider using a proof by contradiction, using Theorem 12.]
6. Let $\overline{AB}, \overline{CD}$ both have midpoint E . Prove that $ACBD$ is a parallelogram.
7. Given points A, B, C such that $AB = AC$, complete a straightedge-compass construction of a rhombus $ABDC$.
8. Prove that, given two distinct points A, B on circle λ which are on the same side of diameter \overline{CD} of λ , that $CB \neq CA$.
9. Given triangle $\triangle ABC$, complete a straightedge-compass construction of a circle that passes through A, B, C . Deduce that given any three points A, B, C that form a triangle (i.e. are not on the same line), there exists a unique circle through these points.
10. Let $\overline{AB}, \overline{CD}$ both have midpoint E and let F, G be points such that $BECF$ and $AEDG$ are parallelograms. Prove that E is the midpoint of FG .