

# Gravity and Electrostatics

- **Newton's Law of Gravity.** Two masses,  $m_1$  and  $m_2$ , experience *gravitational attractive force* to each other, that depends on distance between them,  $r$ :

$$F = -\frac{Gm_1m_2}{r^2}; \quad G = 6.7 \times 10^{-11} \frac{m^3}{kg \cdot s^2}$$

$G$  is called Gravitational Constant. In this equation, '-' sign stands for attraction (positive direction is "away").

- **Coulomb's Law.** Two electric charges,  $q_1$  and  $q_2$ , at distance  $r$ , act onto each other with *electrostatic force* given by Coulomb's formula:

$$F = \frac{kq_1q_2}{r^2}; \quad k = 9 \cdot 10^9 \frac{Nm^2}{C^2}$$

Here  $k$  is called Coulomb's constant. SI unit of electric charge is 1 Coulomb (1C), which is a very large charge. Coulomb's Law is very similar to Newton's, but

- Electric charges can be positive or negative, unlike masses.
- Note that the signs in two laws are different. As a result, charges of the same sign repel, while the opposite ones attract each other.

# Fields

- A **Field** is a physical quantity that has certain value at any point of physical space (x,y,z), and time, t. In other words, it's a function defined in physical space & time.
- A field can be vector or scalar, but there are also other types.
- Electric field **E**, and Newtonian gravity **g**, are examples of vector fields.
- Electric force acting on a charge q:

$$\vec{F}_{elect} = q\vec{E}$$

here electric field does not depend on the charge q itself, but depends on other charges in space.

- Gravitational force:

$$\vec{F}_{grav} = m\vec{g}$$

Mass m is the gravitational charge, g is the local gravity field. g is also an acceleration of a freely falling object, but of course it does not have to have the familiar value of 9.8 m/s<sup>2</sup>, as on the surface of Earth.

# Homework 1

An electric capacitor is made of two parallel metallic plates that are oppositely charged. Let electric field inside this capacitor be  $E=1000\text{N/C}$  (Newton per Coulomb), directed from the positive to negative electrode. When the negative plate is illuminated with UV light, electrons may escape the metal. At moment  $t=0\text{s}$  one such electron appears right near the negative surface, with no initial velocity.

What will be its acceleration, if electron charge is  $e=-1.6 \times 10^{-19} \text{ C}$ , and mass is  $m=0.9 \times 10^{-30}\text{kg}$ .

Calculate the time it will take this electron to reach the positive plate, due to electric force. Distance between the plates is  $h=0.001\text{m}$ .

