

PART 3 Clouds





- Jet streams are fast flowing, relatively narrow air currents found in the atmosphere of some planets, including Earth.
- Discovered in 1940s.
- Can be found in the upper troposphere at ~10-15 km altitude.
- Caused by a <u>combination of the</u> <u>Earth's rotation</u> on its axis and <u>uneven atmospheric heating</u>.
- Strong, high speed (~50-100 mph).
- Major jets move west to east:
 - Polar (strongest)
 - Subtropical



• The polar and subtropical jets merge at some locations and times, while at other times they are well separated.

Jet Streams Role

The path of jet streams <u>steers cyclonic storm systems</u> at lower levels in the atmosphere.



 Jet streams develop meanders, that eventually cut off, detaching and moving air masses.



 In air travel, <u>flight time</u> can be dramatically affected by either flying with the flow or against the flow of a jet stream.

Clouds

A <u>cloud</u> is a visible mass of billions of tiny water droplets or ice crystals suspended in the atmosphere.

- In general, clouds develop in any air mass that becomes saturated (*relative humidity* becomes 100%).
- Saturation occurs due to either or both of two processes: cooling of the air and adding water vapor.





- Ingredients required for cloud formation:
 - water vapor (water in a gaseous state)
 - cooling conditions
 - a surface to condense/deposit on (condensation nuclei)

Condensation Nuclei

Similar to *dew* and *frost*, water vapor <u>requires a surface of</u> <u>some sort to condense on</u> - we call these airborne particles cloud condensation nuclei.



- Terrestrial Sources
 - Dust/sand/dirt particles
 - Smoke volcanic, fires, and pollution
 - Pollens and spores
- Oceanic Sources
 - Sea Salts
- Typical size ~1-2 μm, but can be as large as 100 μm.
 Without these particles, clouds would not form!

Cooling Conditions

Generally, <u>the air must rise</u> in order for it to cool. What can cause the air to rise?

Convectional lifting is associated with heating of the air at the ground surface. This process is active in the interior of continents and near the equator.



Copyright © 2005 Pearson Prentice Hall, Inc.

Orographic uplift occurs when air is forced to rise because of the physical presence of elevated land such as mountains.



Cooling Conditions

rising air

rapid surface cooling

Frontal lifting (convergence) takes place when two masses of air come together. This mechanism of cloud formation is common at mid-latitudes.



Radiative cooling occurs when the Sun is no longer supplying the ground and overlying air with energy (that is, night time). The clouds that result from this type of cooling take the form of surface fog.



Heat radiating from the surface at night, cools the bottom air until it reaches saturation Fog forms first at the surface, thickening as cooling continues.

Basic Cloud Formation

As the air rises, it encounters less pressure. The air parcels expand and cool (called *adiabatic cooling*).



- When the air is cooled to the dew point, <u>condensation</u> <u>occurs</u> and clouds begin to form.
- At higher altitudes, the dew point is the frost point, so water vapor deposition occurs resulting in ice clouds formation.

Floating Conditions



• A cloud will float as long as the drag force of the air dominates over the gravitational force for its constituent water droplets.

- Clouds exist in the <u>moving air</u>: rising up due to *pressure difference* or pushed along by wind.
- Moving air creates an upward <u>drag force</u>.



Precipitation

Droplets suspended in the air will interact with each other, either by colliding and bouncing off each other or by combining to form a larger droplet.

 Eventually, they become large enough so that the acceleration due to gravity is much larger than the acceleration due to drag.





• These relatively large droplets than fall to the ground as precipitation.

Clouds Classification



© 2007 Thomson Higher Education

<u>Based on</u>: • altitude of their bases/bottoms (*alto, strato*) • shape (*cumulus, cirrus*) • presence of rain (*nimbus*)



Lenticular

Mammatus

Nacreous

Kelvin-Helmholtz



