

MATH 7
ASSIGNMENT 12: PASCAL'S TRIANGLE CONTINUED
 JAN 12, 2020

Pascal triangle

Recall the Pascal triangle:

$$\begin{array}{cccccccc}
 & & & & 1 & & & & \\
 & & & & 1 & & 1 & & \\
 & & & 1 & 2 & & 1 & & \\
 & & 1 & 3 & 3 & & 1 & & \\
 & 1 & 4 & 6 & 4 & & 1 & & \\
 1 & 5 & 10 & 10 & 5 & & 1 & & \\
 1 & 6 & 15 & 20 & 15 & 6 & 1 & & \\
 1 & 7 & 21 & 35 & 35 & 21 & 7 & 1 & \\
 1 & 8 & 28 & 56 & 70 & 56 & 28 & 8 & 1
 \end{array}$$

Every entry in this triangle is obtained as the sum of two entries above it. The k -th entry in n -th line is denoted by $\binom{n}{k}$, or by $\binom{n}{k}$. Note that both n and k are counted from 0, not from 1: for example, $\binom{6}{2} = 15$.

These numbers appear in many problems:

- $\binom{n}{k}$ = The number of paths on the chessboard going k units up and $n - k$ to the right
- = The number of words that can be written using k zeros and $n - k$ ones
- = **The number of ways to choose k items out of n (order doesn't matter)**

Principle of Counting

In how many ways can you arrange N objects in a sequence (from the 1st to the N th)?

There are N ways to choose the first one, $N - 1$ ways to choose the second object once the first one has been chosen, $N - 2$ ways to choose the third object once the first and the second objects have been chosen, and so on. So the Answer is $N!$, which is defined by

$$N! := N(N - 1) \dots 1.$$

This is also called P_N , the number of permutations of N elements.

We will see that we can generalize this idea to solve many problems of combinatorics. This idea is sometimes called the *fundamental principle of counting*.

Formula for binomial coefficients

It turns out that there is an explicit formula for $\binom{n}{k}$:

$$\binom{n}{k} = \frac{n(n - 1) \dots (n - k + 1)}{k!} = \frac{n!}{(n - k)!k!}$$

Compare it with the number of ways of choosing k items out of n when the order matters:

$${}_n P_k = n(n - 1) \dots (n - k + 1) = \frac{n!}{(n - k)!}$$

For example, there are $5 \cdot 4 = 20$ ways to choose 2 items out of 5 if the order matters, and $\frac{5 \cdot 4}{2} = 10$ if the order doesn't matter.

Homework

1. How many “words” of length 5 one can write using only letters U and R, namely 3 U’s and 2 R’s? What if you have 5 U’s and 3 R’s? [Hint: each such “word” can describe a path on the chessboard, U for up and R for right. . .]
2. How many sequences of 0 and 1 of length 10 are there? sequences of length 10 containing exactly 4 ones? exactly 6 ones?
3. If we toss a coin 10 times, what is the probability that all will be heads? that there will be exactly one tails? 2 tails? exactly 5 tails?
4. If you have a group of 4 people, and you need to choose one one to go to a competition, how many ways are there to do it? if you need to choose 2? if you need to choose 3?
5. How many ways are there to select 5 students from a group of 12?
6. In a meeting of 25 people, each much shake hands with each other. How many handshakes are there altogether?
7. (a) An artist has 12 paintings. He needs to choose 4 paintings to include in an art show. How many ways are there of doing this?
(b) The same artist now needs to choose 4 paintings to include in a catalog. How many ways are there to do this? (In the catalog, unlike the show, the order matters).
8. (a) There are 15 students in a soccer club. The coach needs to select 11 of them to form the team for a match against another club. How many possibilities does he have?
(b) There are 15 students in a soccer club. The coach needs to select a goalkeeper and 10 players to form the team for a match against another club. How many possibilities does he have?
(The difference between two parts is that in the first case, the coach needs to select 11 players — no need to specify their positions. In the second part, he needs to select 11 players and specify which of them will be the goalkeeper.)

Extra Problems (Optional)

1. A drunkard is walking along a road from the pub to his house, which is located 1 mile north of the pub. Every step he makes can be either to the north, taking him closer to home, or to the south, back to the pub — and it is completely random: every step with can be north of south, with equal chances. What is the probability that after 10 steps, he will move
 - (a) 10 steps north
 - (b) 10 steps south
 - (c) 4 steps north
 - (d) will come back to the starting position
- *2 In a party there are 2020 people, which divide into 505 families, each one composed of a father, a mother, and two children. In the party, there are 505 tables, each one with 4 chairs. At a certain point, all the people in the party took a seat randomly, and it was noticed that each table had exactly a father, a mother and two children, though not necessarily from the same family. What is the probability that each table has the members of the same family?