

## MATH 8: ASSIGNMENT 17

FEB 2ND, 2020

### CIRCLES

Given a circle  $\lambda$  with center  $O$ ,

- A radius is any line segment from  $O$  to a point  $A$  on  $\lambda$ ,
- A chord is any line segment between distinct points  $A, B$  on  $\lambda$ ,
- A diameter is a chord that passes through  $O$ ,
- A tangent line is a line that intersects the circle exactly once; if the intersection point is  $A$ , the tangent is said to be the tangent through  $A$ .

Moreover, we say that two circles are tangent if they intersect at exactly one point.

**Theorem 20.** *Let  $A$  be a point on circle  $\lambda$  centered at  $O$ , and  $m$  a line through  $A$ . Then  $m$  is tangent to  $\lambda$  if and only if  $m \perp \overline{OA}$ . Moreover, there is exactly one tangent to  $\lambda$  at  $A$ .*

*Proof.* First we prove  $(m \text{ is tangent to } \lambda) \implies (m \perp \overline{OA})$ . Suppose  $m$  is tangent to  $\lambda$  at  $A$  but not perpendicular to  $\overline{OA}$ . Let  $\overline{OB}$  be the perpendicular to  $m$  through  $O$ , with  $B$  on  $m$ . Construct point  $C$  on  $m$  such that  $BA = BC$ ; then we have that  $\triangle OBA \cong \triangle OBC$  by *SAS*, using  $OB = OB$ ,  $\angle OBA = \angle OBC = 90^\circ$ , and  $BA = BC$ . Therefore  $OC = OA$  and hence  $C$  is on  $\lambda$ . But this means that  $m$  intersects  $\lambda$  at two points, which is a contradiction.

Now we prove  $(m \perp \overline{OA}) \implies (m \text{ is tangent to } \lambda)$ . Suppose  $m$  passes through  $A$  on  $\lambda$  such that  $m \perp \overline{OA}$ . If  $m$  also passed through  $B$  on  $\lambda$ , then  $\triangle AOB$  would be an isosceles triangle since  $\overline{AO}, \overline{BO}$  are radii of  $\lambda$ . Therefore  $\angle ABO = \angle BAO = 90^\circ$ , i.e.  $\triangle AOB$  is a triangle with two right angles, which is a contradiction.  $\square$

Notice that, given point  $O$  and line  $m$ , the perpendicular  $\overline{OA}$  from  $O$  to  $m$  (with  $A$  on  $m$ ) is the shortest distance from  $O$  to  $m$ , therefore the locus of points of distance exactly  $OA$  from  $O$  should line entirely on one side of  $m$ . This is essentially the idea of the above proof.


**Theorem 21.** *Let  $\overline{AB}$  be a chord of circle  $\lambda$  with center  $O$ . Then  $O$  lies on the perpendicular bisector of  $\overline{AB}$ . Moreover, if  $C$  is on  $\overline{AB}$ , then  $C$  bisects  $\overline{AB}$  if and only if  $\overline{OC} \perp \overline{AB}$ .*

*Proof.* Let  $m$  be the perpendicular bisector of  $\overline{AB}$ . The center  $O$  of  $\lambda$  is equidistant from  $A, B$  by the definition of a circle, therefore by Theorem 14,  $O$  must be on  $m$ . Let  $m$  intersect  $\overline{AB}$  at  $D$ . We then have that  $D$  is the midpoint of  $\overline{AB}$  and also the foot of the perpendicular from  $O$  to  $\overline{AB}$ .

Then if  $C$  bisects  $\overline{AB}$ ,  $C$  lies on the perpendicular bisector  $m$  of  $\overline{AB}$ , which passes through  $O$ , thus  $\overline{OC} \perp \overline{AB}$ . Lastly if  $\overline{OC} \perp \overline{AB}$ , then because there is only one perpendicular to  $\overline{AB}$  through  $O$ , we must have  $C = D$  and hence  $C$  is the midpoint of  $\overline{AB}$ .  $\square$

**Theorem 22.** *Let  $\lambda, \omega$  be circles that intersect at points  $A, B$ . Then  $\overline{AB} \perp \overline{OP}$ .*

*Proof.* We have that  $\triangle AOB$  and  $\triangle APB$  are both isosceles, thus their altitudes from  $O$  and  $P$  respectively both intersect  $\overline{AB}$  at the midpoint  $C$  of  $\overline{AB}$ . Then, since  $m\angle OCA = m\angle ACP = 90^\circ$ , we have that  $m\angle OCP = 180^\circ$ , i.e.  $C$  lies on the line  $\overline{OP}$ . Since  $C$  is the foot of altitudes from  $O$  and  $P$ , this completes the proof.  $\square$




**Theorem 23.** *Let  $\lambda, \omega$  be circles that are both tangent to line  $m$  at point  $A$ . Then  $\lambda, \omega$  are tangent circles.*

*Proof.* Suppose, by contradiction, that  $\lambda, \omega$  intersect at point  $B \neq A$ . Then  $\overline{AB} \perp \overline{OP}$ , therefore both  $\overline{AB}$  and  $m$  are perpendicular to  $\overline{OA}$  through  $A$ . We must therefore have that  $B$  is on  $m$ , but  $m$  is tangent to  $\lambda$  through  $A$ , thus has only intersection with  $\lambda$ , which is a contradiction.  $\square$

## ARCS AND ANGLES

Consider a circle  $\lambda$  with center  $O$ , and an angle formed by two rays from  $O$ . Then these two rays intersect the circle at points  $A, B$ , and the portion of the circle contained inside this angle is called the **arc subtended** by  $\angle AOB$ .

**Theorem 24.** *Let  $A, B, C$  be on circle  $\lambda$  with center  $O$ . Then  $\angle ACB = \frac{1}{2}\angle AOB$ . The angle  $\angle ACB$  is said to be inscribed in  $\lambda$ .*



*Proof.* There are actually a few cases to consider here, since  $C$  may be positioned such that  $O$  is inside, outside, or on the angle  $\angle ACB$ . We will prove the first case here, which is pictured on the left.

*Case 1.* Draw in segment  $\overline{OC}$  and notice that  $m\angle AOC + m\angle BOC + m\angle AOB = 360^\circ$ . Since  $\overline{OC}$  is a radius of  $\lambda$ , we have that  $\triangle AOC$  and  $\triangle BOC$  are isosceles triangles, thus  $m\angle AOC = 180^\circ - 2m\angle OCA$  and  $m\angle BOC = 180^\circ - 2m\angle OCB$ . Therefore we get  $180^\circ - 2m\angle ACO + 180^\circ - 2m\angle BCO + m\angle AOB = 360^\circ \implies m\angle AOB = 2(m\angle ACO + m\angle BCO) \implies m\angle AOB = 2m\angle ACB$ .  $\square$

As a result of Theorem 24, we get that any triangle  $\triangle ABC$  on  $\lambda$  where  $\overline{AB}$  is a diameter must be a right triangle, since the angle  $\angle ACB$  has half the measure of angle  $\angle AOB$ , which is  $180^\circ$ .

The idea captured by the concept of an arc and Theorem 24 is that there is a fundamental relationship between angles and arcs of circles, and that the angle  $360^\circ$  can be thought of as a full circle around a point.

## HOMEWORK

1. Prove that, given a segment  $\overline{AB}$ , there is a unique circle with diameter  $\overline{AB}$ .
2. Given lines  $\overleftrightarrow{AB} \parallel \overleftrightarrow{CD}$  such that  $\overline{AD}, \overline{BC}$  intersect at  $E$  and  $AE = ED$ , prove that  $BE = EC$ .
3. Prove that if a diameter of circle  $\lambda$  is a radius of circle  $\omega$ , then  $\lambda, \omega$  are tangent.
4. Complete the proof of Theorem 24 by proving the cases where  $O$  is not inside the angle  $\angle ACB$ . [Hint: for one of the cases, you may need to write  $\angle ACB$  as the difference of two angles.]
5. Prove the converse of Theorem 24: namely, if  $\lambda$  is a circle centered at  $O$  and  $A, B$ , are on  $\lambda$ , and there is a point  $C$  such that  $m\angle ACB = \frac{1}{2}m\angle AOB$ , then  $C$  lies on  $\lambda$ . [Hint: we need to prove that  $OC = OA$ ; consider using a proof by contradiction, using Theorem 12.]
6. Let  $\overline{AB}, \overline{CD}$  both have midpoint  $E$ . Prove that  $ACBD$  is a parallelogram.
7. Given points  $A, B, C$  such that  $AB = AC$ , complete a straightedge-compass construction of a rhombus  $ABDC$ .
8. Prove that, given two distinct points  $A, B$  on circle  $\lambda$  which are on the same side of diameter  $\overline{CD}$  of  $\lambda$ , that  $CB \neq CA$ .
9. Given triangle  $\triangle ABC$ , complete a straightedge-compass construction of a circle that passes through  $A, B, C$ . Deduce that given any three points  $A, B, C$  that form a triangle (i.e. are not on the same line), there exists a unique circle through these points.
10. Let  $\overline{AB}, \overline{CD}$  both have midpoint  $E$  and let  $F, G$  be points such that  $BECF$  and  $AEDG$  are parallelograms. Prove that  $E$  is the midpoint of  $FG$ .