


Statics

Consider a rigid object in equilibrium. Since it has zero acceleration, and no rotation, the sum of all forces and torques applied should be 0:


$$\sum \vec{F} = 0$$

$$\sum \vec{T} = 0$$



$$\text{Torque } T = F (\text{Force}) \times L (\text{Length})$$

Example: whiteboard on two holders:



We can choose any “axis of rotation” and calculate the torque with respect to it.

- With respect to point P :

$$T_P = mgl_1 - N_2(l_1 + l_2) = 0$$

Note that “clock-wise” is positive direction.

Solving Eq., we obtain: $N_2 = \frac{mgl_1}{l_1 + l_2}$

- Similarly, with respect to point O :


$$T_O = -mgl_2 + N_1(l_1 + l_2) = 0$$

$$N_1 = \frac{mgl_2}{l_1 + l_2} \text{ note that } N_1 + N_2 = mg$$

Homework

Problem 1

- a) A ruler is used to balance two weights as shown in figure 1. The ruler total length is 30 cm, it is supported at its center (at 15 cm mark). Mass $M_1=30$ g, is located at 10 cm mark. The other mass, M_2 is at 30 cm mark. Find M_2
- b) Now the mass M_1 is moved to 0 cm mark. M_2 is still at 30 cm. After that, you can balance the ruler with both masses by placing the support at 10 cm mark. Find the mass of the ruler. Hint: imaging all mass of the ruler to be concentrated at its center, don't try to "break" ruler onto two parts – it's a hard way



Problem 2 (experimental) Use a ruler and a pencil to find the ratio of masses of US quarter and US penny. You may use other two coins if you wish, It's OK to use several identical coins. Make a picture of your experiment, describe procedure and give your results.