

## MATH 5: HANDOUT 21 GEOMETRY 2.

### Sum of angles of an $n$ -gon

Recall that sum of angles of a triangle is  $180^\circ$ . Since a quadrilateral can be cut into 2 triangles, sum of angles of a quadrilateral is  $2 \times 180^\circ = 360^\circ$ . Similarly, for a pentagon we get  $3 \times 180^\circ$ , and for an  $n$ -gon, the sum of angles is  $(n - 2) \times 180^\circ$ .

### Congruence

In general, two figures are called **congruent** if they have same shape and size. We use symbol  $\cong$  for denoting congruent figures: to say that  $M_1$  is congruent to  $M_2$ , we write  $M_1 \cong M_2$ .

Precise definition of what “same shape and size” means depends on the figure. Most importantly, for triangles it means that the corresponding sides are equal and corresponding angles are equal:  $\triangle ABC \cong \triangle A'B'C'$  is the same as:

$$AB = A'B', BC = B'C', AC = A'C', \\ \angle A = \angle A', \angle B = \angle B', \angle C = \angle C'.$$

Note that for triangles, the notation  $\triangle ABC \cong \triangle A'B'C'$  not only tells that these two triangles are congruent, but also shows which vertex of the first triangle corresponds to which vertex of the second one. For example,  $\triangle ABC \cong \triangle PQR$  is not the same as  $\triangle ABC \cong \triangle QPR$ .

### Congruence tests for triangles

By definition, to check that two triangles are congruent, we need to check that corresponding angles are equal and corresponding sides are equal; thus, we need to check 6 equalities. However, it turns out that in fact, we can do with fewer checks.

**Congruence test 1** (Side-Side-Side rule). *If  $AB = A'B'$ ,  $BC = B'C'$  and  $AC = A'C'$  then  $\triangle ABC \cong \triangle A'B'C'$ .*


This rule is commonly referred to as SSS rule.

One can also try other ways to define a triangle by three pieces of information, such as two sides and an angle between them. We will discuss it next time.

This rule — and congruent triangles in general — are very useful for proving various properties of geometric figures. As an illustration, we prove the following useful result.

**Theorem.** *Let  $ABCD$  be a quadrilateral in which opposite sides are equal:  $AB = CD$ ,  $AD = BC$ . Then  $ABCD$  is a parallelogram.*

*Proof.* Let us draw diagonal  $BD$ . Then triangles  $\triangle ABD$  and  $\triangle CDB$  are congruent by SSS; thus, two angles the two angles labeled by letter  $a$  in the figure are equal; also, two angles labeled by letter  $b$  are also equal. Thus, lines  $BC$  and  $AD$  are parallel (alternate interior angles!). In the same way we can show that lines  $AB$  and  $CD$  are parallel. Thus,  $ABCD$  is a parallelogram.  $\square$



(It is also true in the opposite direction: in a parallelogram, opposite sides are equal.)


## Homework

1. Let  $CD$  be a continuation of side  $AC$  in a triangle  $\triangle ABC$ . Show that then  $\angle BCD = \angle A + \angle B$  (such an angle is sometimes called an *exterior angle* of the triangle. [Hint: sum of the angles in a triangle is equal to  $180^\circ$ .])

2. An  $n$ -gon is called *regular* if all sides are equal and all angles are also equal.


- (a) How large is each angle in a regular hexagon (6-gon)?
- (b) Show that in a regular hexagon, opposite sides are parallel. (This is the reason why this shape is used for nuts and bolts).

[Hint: show that each of the angles labeled by letter  $a$  in the figure is equal to  $60^\circ$ , and then use theorem about alternate interior angles.]




3. Let  $ABC$  be a triangle in which two sides are equal:  $AB = BC$  (such a triangle is called *isosceles*). Let  $M$  be the midpoint of the side  $AC$ , i.e.  $AM = MC$ .

- (a) Show that triangles  $\triangle ABM$  and  $\triangle CBM$  are congruent.
- (b) Show that angles  $\angle A$  and  $\angle C$  are equal
- (c) Show that  $\angle AMB = 90^\circ$  (hint:  $\angle AMB = \angle CMB$ ).



4. Let  $ABCD$  be a quadrilateral such that  $AB = BC = CD = AD$  (such a quadrilateral is called *rhombus*). Let  $M$  be the intersection point of  $AC$  and  $BD$ .

- (a) Show that  $\triangle ABC \cong \triangle ADC$



5. In a group of 100 students, 28 speak Spanish, 30 speak German, 42 speak French; 8 students speak Spanish and German, 10 speak Spanish and French, 5 speak German and French and 3 students speak all 3 languages. How many students do not speak any one of the three languages?

[Note: when it says that 28 students speak Spanish, this includes the 8 who speak Spanish and German; similarly for all other combinations.]