Math 4b. Classwork 7. Fractions addition, subtraction.



Addition of fractions with the same denominator.

Let's have a look at the example:


$$\frac{2}{7} + \frac{3}{7} = \frac{5}{7}$$

We divided a whole into 7 small equal parts, took 2 of small parts and then took 3. The result is 5 of $\frac{1}{7}$ parts of a whole. If denominators are different, it's different because parts are not the same size anymore.

Addition and subtraction of fractions with unlike denominators.

Let's try to add $\frac{2}{9}$ and $\frac{2}{3}$. What should we do? What part of the whole the result is?

$$\frac{2}{9} + \frac{1}{3}$$




We divided a whole into 9 small equal parts, took 2 of them. We also divided a whole into 3 bigger parts and took 1 of them. How can we add 2 smaller parts and 1 bigger part? To be able to add parts we need them to be of the same size. We can split each big part into 3 smaller parts.


Now we can add 2 parts and 3 parts of the same size.

$$\frac{2}{9} + \frac{1}{3} = \frac{2}{9} + \frac{1 \cdot 3}{3 \cdot 3} = \frac{2}{9} + \frac{3}{9} = \frac{5}{9}$$

Another example, how to add $\frac{2}{5}$ and $\frac{1}{3}$.




Which part of the whole circle will be the sum?



We need to divide each circle into parts, such that each part will fit into $\frac{2}{5}$ of the circle and $\frac{1}{3}$ of the circle whole number of times. We can divide the circle into 15 parts, 30 parts, 45 parts, etc. All these are common multiples of both denominators 5 and 3. There are many common multiples of 2 numbers, however the most convenient multiple is the Least Common multiple (LCM). LCM (5, 3) = 15. If we split the circle into 15 parts, each such part will fit 6 times into $\frac{2}{5}$ part of the circle, and 5 times into $\frac{1}{3}$ part of the circle.

$$\frac{2}{5} + \frac{1}{3} = \frac{2 \cdot 3}{5 \cdot 3} + \frac{1 \cdot 5}{3 \cdot 5} = \frac{6}{15} + \frac{5}{15} = \frac{11}{15}$$




One more example:

$$\frac{1}{6} + \frac{4}{9} =$$

The most convenient common denominator is 18, LCM of 6 and 9.

$$\frac{1}{6} + \frac{4}{9} = \frac{1 \cdot 3}{6 \cdot 3} + \frac{4 \cdot 2}{9 \cdot 2} = \frac{3}{18} + \frac{8}{18} = \frac{11}{18}$$



When we are talking about fraction, we usually mean the part of a unit as shown in a number line above. **Proper** fractions are parts of a unit; **improper fractions** are the sums of a natural number and a proper fraction. Sometimes we want to find a part of something which is not 1, but can be considered as a single object. For example, among my 30 pencils $\frac{2}{5}$ are yellow. How many yellow pencils do I have? What does it mean to find $\frac{2}{5}$ out of 30? The whole pile of all of all these pencils is a single object and we want to calculate how many pencils does a little pile of $\frac{2}{5}$ of 30 contain? $\frac{2}{5}$ is 2 times $\frac{1}{5}$, and $\frac{1}{5}$ of 30 is 30 ÷ 5. So $\frac{2}{5}$ of 30 pencils will be twice more: $\frac{2}{5} \times 30 = 30 \div 5 \times 2$

Homework.

1. Bring the following fractions to denominator 36, if possible:

$$\frac{7}{12}$$
; $\frac{7}{11}$; $\frac{7}{10}$; $\frac{7}{9}$; $\frac{7}{8}$; $\frac{7}{7}$;

2. Calculate:

$$\frac{1}{5} + \frac{1}{5} + \frac{1}{5} = \qquad \qquad \frac{2}{7} + \frac{1}{7} = \qquad \qquad \frac{7}{9} - \frac{3}{9} =$$

$$\frac{1}{8} + \frac{1}{4} = \qquad \qquad \frac{3}{5} + \frac{2}{6} =$$

3. Simplify the following fractions:

Examples:

$$\frac{3 \cdot 5 \cdot 7}{5 \cdot 7 \cdot 11} = \frac{3}{11}; \qquad \frac{56}{64} = \frac{7 \cdot 8}{8 \cdot 8} = \frac{7}{8}$$

$$\frac{2\cdot 3}{7\cdot 2}; \quad \frac{5\cdot 4}{4\cdot 9}; \quad \frac{7\cdot 5}{2\cdot 7}; \quad \frac{2\cdot 3}{4\cdot 5}$$

$$\frac{22}{66}$$
; $\frac{125}{75}$; $\frac{24}{360}$; $\frac{100}{250}$;

4. Evaluate:

Example:

$$\frac{1}{2} - \frac{1}{3} + \frac{1}{4} = \frac{6}{12} - \frac{4}{12} + \frac{3}{12} = \frac{6 - 4 + 3}{12} = \frac{5}{12}$$

$$\frac{1}{2} - \frac{1}{4} + \frac{3}{5} =$$

$$\frac{3}{4} - \frac{1}{2} + \frac{7}{8} =$$

$$\frac{5}{6} - \frac{2}{3} + \frac{1}{4} =$$

- 5. Math class lasts for $\frac{3}{4}$ of an hour. How many minutes does the class last?
- 6. Julia's father's step is 70 cm long, Julia's step is 20 cm smaller. They start walking making their first step simultaneously. How far they should go to have next simultaneous step?



- 7. Peter spent 2 hours doing his homework. $\frac{1}{3}$ of this time, he spent doing his math homework and $\frac{1}{4}$ of the remaining time he spent on the history assignment. How many minutes did Peter spent on his history assignment and how many minutes did he spent doing his math homework?
- 8. Copy the figure:

