Dividing into groups of equal size.

| 1 Simplify and solve for X. |
|--|
| X - (5 + 3) = 12 |
| X - (6 + 2) = 4 |
| X + 2 - (3 + 6) = 14 |
| X - (3 - 2 + 6) = 10 |
| X + (8 - 4) + 12 - 6 = 20 |
| X + (8 - 4 + 12) - 2 = 16 |
| 2 Compare if possible, using >, <, or = signs (c, x, p, $q \neq 0$). |
| $2 \times c + c \square c \times 3 \qquad \qquad 3 \times c \qquad \square c \times 4 \qquad c \times 6 \square c \times 3 + c \times 2$ |
| $x \times 5 - x \times 2$ $x \times 3$ $p + p \times 2$ $p \times 4$ $q \times 4$ $q + q + q$ |
| 3 Solve the following word problems: |
| A. Barn the dog eats 2 sausages a day. How many sausages will he eat in 7 days? |
| B. Barn the dog eats 2 sausages a day. How many sausages will he eat in x days? |
| C. Barn the dog eats m sausages a day. How many sausages will he eat in x days? |
| D. Barn the dog eats 2 sausages a day. How long will it take him to eat 12 sausages? |
| E. Barn the dog eats 2 sausages a day. How long will it take him to eat x sausages? |
| F. Barn the dog eats w sausages a day. How long will it take him to eat x sausages? |

Lesson 22

Dividing into groups of equal size.

Compare the two word problems and their solution: 4 Problem 2. Foxy Tail wants to treat his Problem I. Foxy Tail wants to treat his friends with apples. He has 16 apples and friends with apples. He has 16 apples intends to give 2 apples to each of his and intends to give the same number of friends. How many friends can he treat apples to each of his two friends. How with apples? many apples will each friend get? 5 Little Joe has decided to share his toy cars among his 5 friends. What is the easiest way to share these cars evenly?

What is the best way to divide these cars into 5 equal groups?

It is easy to split the job in two parts: first, divide the toy cars into groups of 5; second, give each friend a car from each group.



6 There is a large bag of candies. Pop Eye wants to share the bag among Little Joe, Foxy Tail, Jake the Mouse, and himself. Is it possible to divide these candies evenly among the brothers without counting the total number of candies in the bag? Is it always possible to divide all candies evenly?



