


## Exponents

The main reason we use exponents is because it's a shorter way to write out big numbers.

**Exponentiation** is a mathematical operation, written as  $a^n$ , involving two numbers, the **base**  $a$  and the **exponent**  $n$ . When  $n$  is a positive integer, an exponent tells us to multiply the base by itself that number of times: *We can say that  $a$  is raised to the power of  $n$ .*

$a^n$  tells you multiply  $a$  by itself  $n$  times:



$$a^n = \underbrace{a \times a \times \dots \times a}_n$$

$4^3$  This tells us to multiply the base 4 by itself 3 times:  $4^3 = 4 \times 4 \times 4$

When exponent  $n$  of the base  $a$  is a negative integer, it tells us to divide 1 by the base that number of times. Or multiply 1 by the base that number of times and take a reciprocal number

$$a^{-n} = \frac{1}{a^n}$$

### Properties of exponent:

If the same base raised to the different power and then multiplied:

$$b^3 \times b^4 = (b \times b \times b) \times (b \times b \times b \times b) = b \times b \times b \times b \times b \times b \times b = b^{3+4} = b^7$$

$$b^n \times b^m = b^{n+m}$$

If the base raised to the power of n then raised again to the power of m:

$$(b^2)^3 = (b \cdot b)^3 = (b \cdot b) \cdot (b \cdot b) \cdot (b \cdot b) = b^{2 \cdot 3} = b^6$$

$$(b^n)^m = b^{n \cdot m}$$

$$b^1 = b;$$

$$b^0 = 1, \text{ for any } b \text{ except } 0.$$

If two different bases raised to the same power, then:

$$(a \cdot b)^3 = (a \cdot b) \cdot (a \cdot b) \cdot (a \cdot b) = a \cdot a \cdot a \cdot b \cdot b \cdot b = a^3 b^3$$

$$(a \cdot b)^n = a^n b^n$$

## ORDER OF OPERATIONS!!!!!!!

1	2	3	4	5	6
<b>P</b>	<b>E</b>	<b>M</b>	<b>D</b>	<b>A</b>	<b>S</b>
Parentheses	Exponents	Multiplication	Division	Addition	Subtraction
<b>(....)</b>	<b>a<sup>2</sup></b>	<b>X</b>	<b>÷</b>	<b>+</b>	<b>-</b>