Math 5b: Classwork 16 Homework #16 is due February 2

**Definition:** A **rational** number is a number that can be in the form p/q where p and q are integers and q is not equal to zero.

Example: 2/3 is a rational number because 3 and 2 are both integers

**Pigeonhole principle** states that if *n* items are put **into m** <u>pigeonholes</u> with n > m, then at least one pigeonhole must contain more than one item.

Theorem: any rational number is a finite or repeating decimal. The way we proved is using **Pigeonhole principle**.

## **Review**

1. Operations with powers:

$$a^{n} = a \cdot a \cdots a \text{ (ntimes)}$$
$$(a \cdot b)^{n} = a^{n} \cdot b^{n}$$
$$a^{m} \cdot a^{n} = a^{m+n};$$
$$a^{m} \div a^{n} = a^{m-n}$$
$$a^{0} = 1$$
$$a^{-n} = \frac{1}{a^{n}}$$

2. We reviewed solving equations and solving rational equations by multiplying both sides of the equation with the denominator, for example.

$$\frac{(x+1)}{3} = 7$$
$$\frac{(x+1)}{3} \times 3 = 7 \times 3$$
$$(x+1) = 21$$
$$x = 20$$

We also revised the *identities*:

$$(a + b)^{2} = a^{2} + 2ab + b^{2}$$
$$(a - b)^{2} = a^{2} - 2ab + b^{2}$$
$$(a + b)(a - b) = a^{2} - b^{2}$$

And *factorizing*:

$$a(b+c) = ab + ac$$

... and used them to solve equations.

We solved equations with exponents:  $a^x = a^c$  and found out that if we have equal bases we need only compare the exponents (powers) to find the unknown: x = c.

So, we need to find a way to rewrite the equations where both sides have the same base.

## Homework

1. Solve the following equations for *x*: (a)  $\frac{5y-12}{3-2y} = 2$ 

(b) 
$$\frac{8-2x}{3x-1} = 3$$
  
(c)  $\frac{3x+a}{2a-5x} = -1$ 

2. Solve the equation:

$$(x-3)^2 - (x-5)(x+5) = 4$$

3. Simplify the fractions using the above identities and factoring rules:

$$(a)\frac{y^2 - 16}{3y + 12} =$$

(b)
$$\frac{a^2 + 10a + 25}{a^2 - 25} =$$
  
(c) $\frac{15z^2 - 9z}{25z^2 - 9} =$ 

4. Consider the sequence 7, 72, 73,  $\dots$  .7n  $\dots$ 

(a) Show that there will be two numbers in this sequence which have the same last two digits. [*Hint: pigeonhole principle!*]

(b) Show that from some moment, the last two digits of numbers in this sequence will start repeating periodically.

- 5. ABCD, below, is a rectangle which is split into 6 smaller ones by 3 parallel lines. Find:
- (a) The area of each rectangle
- (b) The sum of the areas of the 6 rectangles
- (c) The total area ABCD
- (d) Compare (b) and (c)



- 6. A worker is earning \$24 for each day he works, but he has to pay back \$6 for each day he takes off. After 30 days he ended up receiving no money. How many days did he work?
- 7. Find *n* for
- (a)  $3^{-n} = 3$
- (b)  $3^{-n} = \frac{1}{3}$
- (c)  $9^{-n} = 81$
- 8. (*from 101 puzzles in thought and logic, by C. R. Wylie*) Clark, Jones, Morgan, and Smith are four men whose occupation are butcher, druggist, grocer, and policeman, though not necessarily in that order.
  - Clark and Jones are neighbors and take turns driving each other to work.
  - Jones makes more money than Morgan.
  - Clark beats Smith regularly at bowling.
  - The butcher always walks to work.
  - The policeman doesn't live near the druggist.
  - The only time the grocer and the policeman ever meet is when the policeman arrested the grocer for speeding.
  - The policeman makes more money than the druggist or the grocer.

What is each man's occupation?