

Math 5b: Classwork 18

Homework #18 is due February 23 (NO CLASS NEXT SUNDAY, Feb.16)

If you did not hand Homework 17 today, please hand it in next time on Feb., 23

Definition: A **rational** number is a number that can be in the form p/q where p and q are integers and q is not equal to zero.

Example: $2/3$ is a rational number because 3 and 2 are both integers

Pigeonhole principle states that if n items are put into m **pigeonholes** (containers) with $n > m$, then at least one pigeonhole (container) must contain more than one item.

In layman's terms, if you have more "objects" than you have "holes," at least one hole must have multiple objects in it.

Theorem: any rational number is a finite or repeating decimal. The way we proved is using **Pigeonhole principle**.

Example of the proof is provided in Problem #8 here -> <https://www.math.ksu.edu/~zlin/m510/hw1.pdf>

Proof. First we assume that n is a positive integer. Apply the division algorithm to get $m = qn + r_0$ with $0 \leq r_0 \leq n - 1$. Here q is the integer part of the rational number m/n . To compute the tenth place digit a_1 , one uses the division algorithm $r_0 \times 10 = a_1 n + r_1$ with $0 \leq r_1 \leq n - 1$. More generally, to get the 10^{-i} place digit a_i , one uses the remainder r_{i-1} and division algorithm $r_{i-1} \times 10 = a_i n + r_i$. When $i = n$, then the $n+1$ remainders r_0, r_1, \dots, r_n have value ranging from 0 to $n-1$. Thus, by the pigeonhole principle, there must be two equal remainders, say, $r_i = r_j$ with $i < j$. Let j be the smallest possible number such that $r_j = r_i$ for some $i < j$. Then $a_{j+1} = a_{i+1}$ with $r_{j+1} = r_{i+1}$ and, inductively, $a_{i+k} = a_{j+k}$ and $a_i = a_{i+(j-i)} = a_{i+2(j-i)} = \dots$. This shows that the decimal is repeating with the repeating part $a_i a_{i+1} \dots a_{j-1}$.

Review

1. Operations with powers:

$$a^n = a \cdot a \cdot \dots \cdot a \text{ (n times)}$$

$$(a \cdot b)^n = a^n \cdot b^n$$

$$a^m \cdot a^n = a^{m+n};$$

$$a^m \div a^n = a^{m-n}$$

$$a^0 = 1$$

$$a^{-n} = \frac{1}{a^n}$$

2. We reviewed solving equations and solving rational equations by multiplying both sides of the equation with the denominator, for example.

$$\frac{(x+1)}{3} = 7$$

$$\frac{(x+1)}{3} \times 3 = 7 \times 3$$

$$(x+1) = 21$$

$$x = 20$$

We also revised the *identities*:

$$(a+b)^2 = a^2 + 2ab + b^2$$

$$(a-b)^2 = a^2 - 2ab + b^2$$

$$(a+b)(a-b) = a^2 - b^2$$

And *factorizing*:


$$a(b+c) = ab + ac$$

... and used them to solve equations.

We solved equations with exponents: $a^x = a^c$ and found out that if we have equal bases we need only compare the exponents (powers) to find the unknown: $x = c$.

So, we need to find a way to rewrite the equations where both sides have the same base.


Geometry: Angles



Opposite angles, formed from crossing straight lines, are equal.

$\angle \alpha = \angle \alpha$ – opposite

$\angle \alpha + \angle \beta = 180^\circ$ – on a straight line,



$l_1 \parallel l_2$ t is transversal:

$$\angle 1 = \angle 2 = \angle 3$$


$\angle 1 = \angle 3$ = alternate internal angles

$\angle 1 = \angle 2$ = corresponding angles

Or complementary angles

$\angle 4 = \angle 2 =$ alternate exterior angles

From both these pieces of information we can show that the sum of angles in a triangle is always 180° .



Homework 18

1. Can one cut from the circle of diameter 9.9 m?

- a) A rectangle with the sides of 6m and 8m;
- b) A square with the side of 7m;

Please provide the step by step solution similar to how we did it in class.

2. The area of an isosceles (2 equal sides -legs) triangle is 60 square meters. The height drawn to the base (third side) is 5m. *The height of an isosceles triangle divides its base side in half. Find all sides of the triangle. Please write out the full solution similar to how it was done in class.*

The area of the triangle is **half of the base times height**.

- "b" is the distance along the base
- "h" is the height (measured at right angles to the base)

$$\text{Area} = \frac{1}{2} \times b \times h$$

$$a(b + c) = ab + ac$$

$$a(b - c) = ab - ac$$

3. Solve the following equations for *(don't forget to check your solutions)*:

$$(a) \frac{12-10x}{1-5x} = 4$$

$$(b) \frac{7x-9}{1-3x} - 8 = 4$$

$$(c) \frac{6y-18}{10y-30} - y = 2$$

$$(d) \frac{35z-14}{4-10z} + 3z = 2$$

$$(e) \frac{5z+b}{2b-3z} - 3 = -1$$

4. Solve the equation *(after reading review section above)*:

$$(x + 7)^2 - (x - 6)(x + 6) = 1$$

5. Simplify the fractions using the above identities and factoring rules *(after reading review section above)*:

$$(a) \frac{4y^2-25}{6y+15} =$$

$$(b) \frac{9a^2+24a+16}{32-18a^2} =$$

$$(c) \frac{56z^2-105z}{225-64z^2} =$$

$$(d) \frac{36-60x+25x^2}{15x-18} =$$

6. Find n for

$$(a) 5^{-2n+3} = 125$$

$$(b) 5^{-8n} = 1/625$$

$$(c) 8^{-5n-7} = 64^9$$

7. Find the following square-roots. For the final answer if you cannot calculate the exact value of the square root, leave it as a square root.

(a) $\sqrt{11^{28}}$

(b) $(\sqrt{11})^{28}$

(c) $\sqrt{11^{28}} / \sqrt{11^{84}}$

(d) $\frac{\sqrt{11^{20}}}{121}$

(e) $\frac{\sqrt{11^{20}}}{121} * \sqrt{11^{15}}$

(f) $\frac{\sqrt{11^{20}}}{121} * \sqrt{11 * 11^{15}}$

(g) $\frac{\sqrt{11^{20}}}{121} * \sqrt{11 * 11^{-15}}$

(h) $\frac{\sqrt{11^{-20}}}{121} * \sqrt{11 * 11^{-17}} / \sqrt{11^{-15-22} / 11^9}$

8. Simplify

(a) $2 / (x + 2) - 2 / (x - 2)$

(b) $(2 + 2 / x) / (x + 1)$

(c) $(2 + 2 / x) / (x + 2)$


(d) $(2 + 2 / x) / (2 - 2 / x)$

9. If $a = 7^{-32}5^{19}$, $b = 21^{35}3^{-43}$, $c = 35^{72}2^{-23}$, and $d = 10^{19}2^{44}3^{-76}$ what is the value of ab ? of a/b ? abc ? ab/c ? $abcd$? ab/cd ? Prime factors only, simplify as much as possible.
10. Using the same sequence of steps we did in class, perform division of $80/7$. Show all A_i (quotient) and R_i (remainders). How many “pigeonholes” do we have in this case and what goes into those pigeonholes?

Homework 17

1. On the picture, a and b , which are parallel to each other, are intersected by line c . What are the relationships:

- (a) $\angle 3$ and $\angle 5$
- (b) $\angle 2$ and $\angle 8$
- (c) Prove that $\angle 4 + \angle 5 = 180^\circ$.



2. In the same picture,


- (a) if $\angle 7 = 65^\circ$, find: $\angle 1$, $\angle 3$, $\angle 1 + \angle 6$
- (b) If you know that $\angle 7 = \angle 1$, prove that*: $\angle 1 = \angle 3$ and $\angle 5 = \angle 1$

(* or say why the angles will be equal)

1) B

3. Intersecting at point B on triangle ABC is drawn line DS, such that DS is parallel to AC. Prove that (or say why the angles will be equal):

- (a) $\angle ACB = \angle SBC$
- (b) $\angle CAB = \angle DBA$
- (c) $\angle CAB = \angle SBK$
- (d) If $\angle CAB = 40^\circ$ and $\angle BCA = 60^\circ$, find angles $\angle ABD$ and $\angle SBC$



4. In triangle ABC, $\angle A = 35^\circ$, $\angle B = 55^\circ$, prove that this triangle is right-angled.
5. What type of triangle has one angle equal to the sum of the other two?
6. Find each of the outside angles of a right-triangle, if one of its angles is 58° .