

Gravity and Electrostatics

- **Newton's Law of Gravity.** Two masses, m_1 and m_2 , experience *gravitational attractive force* to each other, that depends on distance between them, r :

$$F = -\frac{Gm_1m_2}{r^2}; \quad G = 6.7 \times 10^{-11} \frac{m^3}{kg \cdot s^2}$$

G is called Gravitational Constant. In this equation, '-' sign stands for attraction (positive direction is "away").

- **Coulomb's Law.** Two electric charges, q_1 and q_2 , at distance r , act onto each other with *electrostatic force* given by Coulomb's formula:

$$F = \frac{kq_1q_2}{r^2}; \quad k = 9 \cdot 10^9 \frac{Nm^2}{C^2}$$

Here k is called Coulomb's constant. SI unit of electric charge is 1 Coulomb (1C), which is a very large charge. Coulomb's Law is very similar to Newton's, but

- Electric charges can be positive or negative, unlike masses.
- Note that the signs in two laws are different. As a result, charges of the same sign repel, while the opposite ones attract each other.

Homework

Problem 1

- a) By using Newton's law of gravity, find the gravitational acceleration on the surface of a planet with mass M and radius R . For doing this, consider an apple of mass m . Its weight is mg . But it also must be equal to Newton's gravitational force.
- b) Imagine that you discovered a planet with the same density as Earth, but its radius is twice as big. What will be the value of g on that planet?

Problem 2

Two identical charges placed at distance **10 cm** from each other experience repulsive force 0.1N. Determine the magnitude of the charges.