

MATH 7: HANDOUT 11
POKER PROBABILITIES

POKER PROBABILITIES

In the game of poker, a player is dealt five cards from a regular deck with 4 suits ($\spadesuit, \clubsuit, \diamondsuit, \heartsuit$) with card values in the following order: A, 2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A. We calculated probabilities of the following combinations:

Royal Flush: 10, J, Q, K, A of any suit (Example: $10\heartsuit, J\heartsuit, Q\heartsuit, K\heartsuit, A\heartsuit$)

There are only 4 of them.

Straight Flush: Five cards in a row of the same suit (Example: $6\heartsuit, 7\heartsuit, 8\heartsuit, 9\heartsuit, 10\heartsuit$)

Each of these can start from any card from A to 9, and be in each of the four suits: $9 \times 4 = 36$. Notice that we excluded royal flushes from our computation (if we start with 10, we get a Royal Flush).

Four of a kind: Four cards of the same value, and one other random card (Example: $K\heartsuit, K\spadesuit, K\diamondsuit, K\clubsuit, 2\clubsuit$)

Which card $13 \times$ Which other value $12 \times$ Which other suit $4 = 13 \cdot 12 \cdot 4$.

Full House: Three cards of the same value, and two cards of the same value (Example: $K\heartsuit, K\spadesuit, K\diamondsuit, 4\spadesuit, 4\clubsuit$)

Which card for 3 $13 \times$ Which three suits $\binom{4}{3} \times$ Which card for a pair $12 \times$ Which two suits $\binom{4}{2} = 13 \binom{4}{3} \cdot 12 \binom{4}{2}$.

Flush: Five cards of the same suit, not in order (Example: $3\heartsuit, 6\heartsuit, 8\heartsuit, J\heartsuit, A\heartsuit$)

Which suit $4 \times$ Which five cards $\binom{13}{5} = 4 \binom{13}{5}$. We also need to exclude Royal Flushes and Straight Flushes, so the total is $4 \binom{13}{5} - 40$.

Straight: Five cards in order, possibly of different suits (Example: $5\heartsuit, 6\spadesuit, 7\diamondsuit, 8\spadesuit, 9\clubsuit$)

Which card to start from (anything from A to 10) $10 \times$ Five suits $4^5 = 10 \cdot 4^5$. From here we also need to exclude Royal Flushes and Straight Flushes, so the final answer is $10 \cdot 4^5 - 40$.

Triple: Three cards of the same value, and two other random cards (Example: $K\heartsuit, K\spadesuit, K\diamondsuit, 4\spadesuit, 2\clubsuit$)

Which card $\binom{13}{1} \times$ Which three suits $\binom{4}{3} \times$ Which two other values $\binom{12}{2} \times$ Which two suits for these two random cards $4^2 = \binom{13}{1} \binom{4}{3} \binom{12}{2} 4^2$.

Two pairs: Two cards of the same value, two cards of the same value, and a random card (Example: $K\heartsuit, K\spadesuit, 10\diamondsuit, 10\spadesuit, 4\clubsuit$)

Pair: Two cards of the same value, and three other random cards (Example: $K\heartsuit, K\spadesuit, Q\diamondsuit, 4\spadesuit, 2\clubsuit$)

To calculate probabilities of each of these combinations, we have to divide the counts above by the total number of poker hands, which is $\binom{52}{5}$. The table below gives the probabilities and odds:

Combination	Count	Probability	Odds
Royal Flush	4	0.000154%	1 : 649,740
Straight Flush	36	0.00139%	1 : 72,192
Four of a Kind	$13 \cdot 12 \cdot 4$	0.024%	1 : 4,165
Full House	$13 \binom{4}{3} \cdot 12 \binom{4}{2}$	0.1441%	1 : 693
Flush	$4 \binom{13}{5} - 40$	0.1965%	1 : 508
Straight	$10 \cdot 4^5 - 40$	0.3925%	1 : 254
Triple	$\binom{13}{1} \binom{4}{3} \binom{12}{2} 4^2$	2.1128%	1 : 46.3
Two Pairs			
Pair			
Nothing			

HOMEWORK

- There is only one problem in this homework.

Note that we did not calculate the probabilities of two poker combinations, **Two Pairs** and **Pair**. Please study this handout in detail and use the same logic as for other combinations to find the probabilities of these combinations. Then fill out the line for **Nothing** – probability that there is a poker hand with nothing interesting in it.