

A good example of such system is the ancient Egyptian decimal numeral system.

						
1	10	100	1000	10000	100000	10^6

Egyptian numeral hieroglyphs



A close-up photograph of a stone inscription. At the top, there is a stylized bull's head (djedw) hieroglyph. Below it are several lines of hieroglyphic text, including a row of three lotus flowers and a row of three ankh symbols.

The example on the right is from the Louvre and has the writing in columns. Notice the calf at the top of the picture is facing to the right. Thus, the hieroglyphs are read from top to bottom and, within each line, from right to left. The number pictured is composed of four lotus flowers (4000), six stacked coils of rope (600), two hobbles (20), and two tally marks (2), namely 4622. Since the number follows “calf,” it translates as 4622 calves.

Author(s): Cynthia J. Huffman (Pittsburg State University)

All photos in this article—except for Figures 2–4, 15, and 40—were taken by the author and can be used with attribution under a [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/) license. <https://maa.org/press/periodicals/convergence/an-ancient-egyptian-mathematical-photo-album-hieroglyph-numerals-and-more>



The Babylonian system (from about 2000 BC) used only two symbols to write any number between 1 and 60, which shows that it was based on 60 and had a somewhat 10-based system inside:

to count units and $\angle$ to count tens.

▽	1	◁▽	11	◁◁▽	21	◁◁◁▽	31	◁◁◁▽	41	◁◁◁▽	51
▽▽	2	◁▽▽	12	◁◁▽▽	22	◁◁◁▽▽	32	◁◁◁▽▽	42	◁◁◁▽▽	52
▽▽▽	3	◁▽▽▽	13	◁◁▽▽▽	23	◁◁◁▽▽▽	33	◁◁◁▽▽▽	43	◁◁◁▽▽▽	53
▽▽▽	4	◁▽▽▽	14	◁◁▽▽▽	24	◁◁◁▽▽▽	34	◁◁◁▽▽▽	44	◁◁◁▽▽▽	54
▽▽▽	5	◁▽▽▽	15	◁◁▽▽▽	25	◁◁◁▽▽▽	35	◁◁◁▽▽▽	45	◁◁◁▽▽▽	55
▽▽▽	6	◁▽▽▽	16	◁◁▽▽▽	26	◁◁◁▽▽▽	36	◁◁◁▽▽▽	46	◁◁◁▽▽▽	56
▽▽▽	7	◁▽▽▽	17	◁◁▽▽▽	27	◁◁◁▽▽▽	37	◁◁◁▽▽▽	47	◁◁◁▽▽▽	57
▽▽▽	8	◁▽▽▽	18	◁◁▽▽▽	28	◁◁◁▽▽▽	38	◁◁◁▽▽▽	48	◁◁◁▽▽▽	58
▽▽▽	9	◁▽▽▽	19	◁◁▽▽▽	29	◁◁◁▽▽▽	39	◁◁◁▽▽▽	49	◁◁◁▽▽▽	59
◁	10	◁◁	20	◁◁◁	30	◁◁◁	40	◁◁◁	50		

[This Photo](#) by Unknown Author is licensed under [CC BY-NC-ND](#)

Also, it was a positional value system. For example, number 62 was shown as $\overline{\text{I}} \text{ II}$, which means one time 60 and 2. The use of this sexagesimal (60-based) system is still noticeable today, with 60 minutes in one hour, 60 seconds in a minute, 360° for the full turn.

Now we adopt a notation where we separate the numerals by commas, when we use our digits to represent Babylonian notation, for example, 1,57,46,40 represents the sexagesimal number

$$60^3 \cdot 1 + 60^2 \cdot 57 + 60^1 \cdot 46 + 60^0 \cdot 40$$

which, in decimal notation is 424000.

$\overline{\text{I}}$	II	IV	X
1,57,46,40 = 424000			

Here is **1,57,46,40 in Babylonian numerals**

https://mathshistory.st-andrews.ac.uk/HistTopics/Babylonian_mathematics/



Babylonian mathematical tablet Plimpton 322.Credit...Christine Proust and Columbia University

Another very well-known numeral system is the Roman system; it was used for thousands of years and, in some cases, is still used today as well. It's a “decimal”, 10-based system, but the symbols (letters)are used in an unusual way.

Symbol	I	V	X	L	C	D	M
Value	1	5	10	50	100	500	1000

1	2	3	4	5	6	7	8	9	10
I	II	III	IV	V	VI	VII	VIII	IX	X

For example, 4 is one less than 5, so 4 can be written as IV. Same principle of subtractive notation is used for 9 -> IX, 40 and 90 -> XL and XC, 400 and 900 -> CD and CM

Some other examples:

- $29 = XX + IX = \mathbf{XXIX}$.
- $347 = CCC + XL + VII = \mathbf{CCCXLVII}$.
- $789 = DCC + LXXX + IX = \mathbf{DCCLXXXIX}$.
- $2,421 = MM + CD + XX + I = \mathbf{MMCDXXI}$

Any missing place (represented by a zero in the place-value equivalent) is omitted, as in Latin (and English) speech:

- $160 = C + LX = \mathbf{CLX}$
- $207 = CC + VII = \mathbf{CCVII}$
- $1,009 = M + IX = \mathbf{MIX}$
- $1,066 = M + LX + VI = \mathbf{MLXVI}$

Can non-decimal place-value system be created? For example, with base 5?

Let see, how we can create this kind of system (we use our normal digits).

Num ₁₀	1	2	3	4	5	6	7	8	9	10
Num ₅	1	2	3	4	10	11	12	13	14	20

11	12	13	14	15	16	17	18	19	20
21	22	23	24	30	31	32	33	34	40

21	22	23	24	25	26	27	28	29	30
41	42	43	44	100	101	102	103	104	105

We only have 5 digits (0, 1, 2, 3, 4), and 4 first “natural” numbers in such system will be represented as one-digit numbers. Number 5 then should be shown as a 2-digit number, with first digit 1 (place - value equal to 5^1) and 0 of “units”. Any number is now written in the form

$$\dots + 5^3 \cdot n + 5^2 \cdot m + 5^1 \cdot k + 5^0 \cdot p, \quad n, m, k, p \text{ are } 0, 1, 2, 3, 4$$

$$33 = 25 + 5 + 3 = 5^2 \cdot 1 + 5^1 \cdot 1 + 5^0 \cdot 3 = 113_5$$

$$195 = 125 + 25 \cdot 2 + 5 \cdot 4 = 5^3 \cdot 1 + 5^2 \cdot 2 + 5^1 \cdot 4 + 0 = 1240_5$$

And vice versa, we can transform the number from 5-base to decimal system:

$$2312_5 = 5^3 \cdot 2 + 5^2 \cdot 3 + 5^1 \cdot 1 + 2 = 250 + 75 + 5 + 2 = 332$$

Let's do the addition of 1240_5 and 2312_5 (I omitted $_5$ notation):

$$\begin{array}{r} 1240 \\ + 2312 \\ \hline 4102 \end{array}$$

$$4102_5 = 5^3 \cdot 4 + 5^2 \cdot 1 + 5^1 \cdot 0 + 2 = 125 \cdot 4 + 25 \cdot 1 + 0 + 2 = 500 + 25 + 2 = 527$$

Let's try to introduce a new digit **S** for 10 and then create an 11 based system.

$$11^2 = 121, \quad 11^3 = 1331$$

$$890 = 121 \cdot 7 + 11 \cdot 3 + 10 = 11^2 \cdot 7 + 11^1 \cdot 3 + 10 = 73S_{11}$$

$$4S2_{11} = 11^2 \cdot 4 + 11^1 \cdot 10 + 2 = 484 + 110 + 2 = 596$$

There is another very important place-value system: binary system, base 2 system where only two digits exist; 0, and 1.

Num ₁₀	1	2	3	4	5	6	7	8	9	10
Num ₂	1	10	11	100	101	110	111	1000	1001	1010

In this system place value of a digit is a power of 2:

$$\dots + 2^3 \cdot (0,1) + 2^2 \cdot (0,1) + 2^1 \cdot (0,1) + 2^0 \cdot (0,1)$$

$$11 = 8 + 2 + 1 = 2^3 \cdot 1 + 2^2 \cdot 0 + 2^1 \cdot 1 + 2^0 \cdot 1 = 1011_2$$

$$75 = 64 + 8 + 2 + 1 = 2^6 \cdot 1 + 2^5 \cdot 0 + 2^4 \cdot 0 + 2^3 \cdot 1 + 2^2 \cdot 0 + 2^1 \cdot 1 + 2^0 \cdot 1 = 1001011_2$$

$$189 = 128 + 32 + 16 + 8 + 4 + 1$$

$$= 2^7 \cdot 1 + 2^6 \cdot 0 + 2^5 \cdot 1 + 2^4 \cdot 1 + 2^3 \cdot 1 + 2^2 \cdot 1 + 2^1 \cdot 0 + 2^0 \cdot 1$$

$$= 10111101_2$$

Binary numbers can be transferred to decimals too. We will do it from right to left for convenience.

$$\begin{aligned} 11010111_2 &= 2^0 \cdot 1 + 2^1 \cdot 1 + 2^2 \cdot 1 + 2^3 \cdot 0 + 2^4 \cdot 1 + 2^5 \cdot 0 + 2^6 \cdot 1 + 2^7 \cdot 1 \\ &= 128 + 64 + 16 + 4 + 2 + 1 = 215 \end{aligned}$$

Exercises:

1. Write numbers 51 and 175 in binary system.
2. Write the numbers, written in the binary system in decimal system:
a. 11011011_2 ; b. 10001101_2 , c. 11111111_2
3. Write the numbers 245 and 324 in 6-based place-value system. Remember, that in this system you will have only 0, 1, 2, 3, 4, and 5 as digits.
4. Write the numbers 234_6 and 403_6 written in the 6-based place-value system (small number 6 shows that the number is not in decimal, but in 6-based system) in decimal system.
5. How to arrange 127 1-dollar bills in seven wallets so that any amount from 1 to 127 dollars could be issued without opening the wallets?
6. Robert thought of a number not less than 1 and not more than 1000. Julia is allowed to ask only such questions to which Robert can answer “yes” or “no” (Robert always tells the truth). Can Julia determine the hidden number in 10 questions?
7. There is a bag of sugar, a scale and a weight of 1 g. Is it possible to measure 1 kg of sugar in 10 weights?